

JEE MAIN 2021

ONLINE 22nd July 2nd Shift

PHYSICS

SECTION-A (MULTIPLE CHOICE QUESTIONS)

1. Match List-I with List-II. List-I

List-I	List-II
(A) $\omega L > \frac{1}{\omega C}$	(i) Current is in phase with emf
(B) $\omega L = \frac{1}{2}$	(ii) Current lags behind

ii) Current lags behind the applied emf

(iii) Maximum current occurs

(C)
$$\omega L < \frac{1}{\omega C}$$

 ωC

(D) Resonant frequency (iv) Current leads the emf Choose the correct answer from the options given below:

- (a) (A) (iv), (B) (iii), (C) (ii), (D) (i)
 (b) (A) (ii), (B) (i), (C) (iv), (D) (iii)
- (c) (A) (iii), (B) (i), (C) (iv), (D) (ii)
- (d) (A) (ii), (B) (i), (C) (iii), (D) (iv)
- 2. Statement-I : The ferromagnetic property depends on temperature. At high temperature, ferromagnet becomes paramagnet.

Statement-II : At high temperature, the domain wall area of a ferromagnetic substance increases.

In the light of the above statements, choose the most appropriate answer from the options given below.

- (a) Both Statement-I and Statement-II are true.
- (b) Both Statement-I and Statement-II are false.
- (c) Statement-I is true but Statement-II is false.
- (d) Statement-I is false but Statement-II is true.
- 3. Consider a situation in which reverse biased current of a particular *P*-*N* junction increases when it is exposed to a light of wavelength ≤ 621 nm. During this process, enhancement in carrier concentration take place due to generation of hole -electron pairs. The value of band gap is

(a)	4 eV	(b)	1 eV
(c)	0.5 eV	(d)	$2 \ eV$

4. A copper (Cu) rod of length 25 cm and cross-sectional area 3 mm² is joined with a similar aluminium (Al) rod as shown in figure. Find the resistance of the combination between the ends *A* and *B*. (Take resistivity of copper = $1.7 \times 10^{-8} \Omega$ m, Resistivity of aluminium = $2.6 \times 10^{-8} \Omega$ m).



(c) $0.0858 \text{ m}\Omega$ (d) $1.420 \text{ m}\Omega$

5. In a circuit consisting of a capacitance and a generator with alternating emf $E_g = E_{g_0} \sin\omega t$, V_C and I_C are the voltage and current. Correct phasor diagram for such circuit is



6. What will be the projection of vector $\vec{A} = \hat{i} + \hat{j} + \hat{k}$ on vector $\vec{B} = \hat{i} + \hat{j}$?

(a)
$$\sqrt{2}(\hat{i}+\hat{j})$$
 (b) $(\hat{i}+\hat{j})$

(c)
$$\sqrt{2} (\hat{i} + \hat{j} + \hat{k})$$
 (d) $2 (\hat{i} + \hat{j} + \hat{k})$

- 7. Consider a situation in which a ring, a solid cylinder and a solid sphere roll down on the same inclined plane without slipping. Assume that they start rolling from rest and having identical diameter. The correct statement for this situation is
 - (a) The ring has the greatest and the cylinder has the least velocity of the centre of mass at the bottom of the inclined plane.
 - (b) The cylinder has the greatest and the sphere has the least velocity of the centre of mass at the bottom of the inclined plane.

- (c) The sphere has the greatest and the ring has the least velocity of the centre of mass at the bottom of the inclined plane.
- (d) All of them will have same velocity.
- 8. What should be the height of transmitting antenna and the population covered if the television telecast is to cover a radius of 150 km? The average population density around the tower is 2000/km² and the value of $R_e = 6.5 \times 10^6$ m.
 - (a) Height = 1800 m Population covered = 1413×10^8
 - (b) Height = 1600 m Population covered = 2×10^5
 - (c) Height = 1731 m Population covered = 1413×10^5
 - (d) Height = 1241 m Population covered = 7×10^5
- 9. Intensity of sunlight is observed as 0.092 Wm⁻² at a point in free space. What will be the peak value of magnetic field at that point? ($\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$) (a) $1.96 \times 10^{-8} \text{ T}$ (b) $2.77 \times 10^{-8} \text{ T}$
 - (a) 1.90×10^{-1} (b) 2.77×10^{-1} (c) 5.88 T (d) 8.31 T
- 10. An electric dipole is placed on *x*-axis in proximity to a line charge of linear charge density 3.0×10^{-6} C/m. Line charge is placed on *z*-axis and positive and negative charge of dipole is at a distance of 10 mm and 12 mm from the origin respectively. If total force of 4 N is exerted on the dipole, find out the amount of positive or negative charge of the dipole.
 - (a) $4.44 \,\mu C$ (b) $8.8 \,\mu C$
 - (c) 0.485 mC (d) 815.1 nC
- 11. A nucleus with mass number 184 initially at rest emits an α -particle. If the *Q* value of the reaction is 5.5 MeV, calculate the kinetic energy of the α -particle.
 - (a) 5.38 MeV (b) 0.12 MeV
 - (c) 5.0 MeV (d) 5.5 MeV
- **12.** A body is projected vertically upwards from the surface of earth with a velocity sufficient enough to carry it to infinity. The time taken by it to reach height *h* is ______s.

(a)
$$\sqrt{\frac{2R_e}{g}} \left[\left(1 + \frac{h}{R_e} \right)^{3/2} - 1 \right]$$

(b) $\frac{1}{3} \sqrt{\frac{R_e}{2g}} \left[\left(1 + \frac{h}{R_e} \right)^{3/2} - 1 \right]$
(c) $\sqrt{\frac{R_e}{2g}} \left[\left(1 + \frac{h}{R_e} \right)^{3/2} - 1 \right]$
(d) $\frac{1}{3} \sqrt{\frac{2R_e}{g}} \left[\left(1 + \frac{h}{R_e} \right)^{3/2} - 1 \right]$

- 13. Choose the correct option.
 - (a) True dip is always greater than the apparent dip.
 - (b) True dip is less than the apparent dip.
 - (c) True dip is not mathematically related to apparent dip.
 - (d) True dip is always equal to apparent dip.
- 14. A ray of light passes from a denser medium to a rarer medium at an angle of incidence *i*. The reflected and refracted rays make an angle of 90° with each other. The angle of reflection and refraction are respectively r and r'. The critical angle is given by



- 15. T_0 is the time period of a simple pendulum at a place. If the length of the pendulum is reduced to $\frac{1}{16}$ times of
 - its initial value, the modified time period is

(a)
$$8\pi T_0$$
 (b) $\frac{1}{4}T_0$

(c)
$$T_0$$
 (d) $4T_0$

- **16.** A bullet of 4 g mass is fired from a gun of mass 4 kg. If the bullet moves with the muzzle speed of 50 ms⁻¹, the impulse imparted to the gun and velocity of recoil of gun are
 - (a) 0.4 kg m s^{-1} , 0.1 m s^{-1} (b) 0.4 kg m s^{-1} , 0.05 m s^{-1}
 - (c) 0.2 kg m s^{-1} , 0.05 m s^{-1} (d) 0.2 kg m s^{-1} , 0.1 m s^{-1}
- 17. The motion of a mass on a spring with spring constant *K* is as shown in figure.



The equation of motion is given by $x(t) = A\sin\omega t + B\cos\omega t$ with $\omega = \sqrt{\frac{K}{m}}$. Suppose that at time t = 0, the position of mass is x(0) and velocity v(0), then its displacement can also be represented as $x(t) = C\cos(\omega t - \phi)$, where *C* and ϕ are

(a)
$$C = \sqrt{\frac{2\nu(0)^2}{\omega^2} + x(0)^2}, \ \phi = \tan^{-1}\left(\frac{x(0)\omega}{2\nu(0)}\right)$$

(b) $C = \sqrt{\frac{\nu(0)^2}{\omega^2} + x(0)^2}, \ \phi = \tan^{-1}\left(\frac{x(0)\omega}{\nu(0)}\right)$



(c)
$$C = \sqrt{\frac{2\nu(0)^2}{\omega^2} + x(0)^2}, \ \phi = \tan^{-1}\left(\frac{\nu(0)}{x(0)\omega}\right)$$

(d) $C = \sqrt{\frac{\nu(0)^2}{\omega^2} + x(0)^2}, \ \phi = \tan^{-1}\left(\frac{\nu(0)}{x(0)\omega}\right)$

18. An electron of mass m_e and a proton of mass m_p are accelerated through the same potential difference. The ratio of the de-Broglie wavelength associated with the electron to that with the proton is

(a) 1 (b)
$$\frac{m_p}{m_e}$$
 (c) $\frac{m_e}{m_p}$ (d) $\sqrt{\frac{m_p}{m_e}}$

- **19.** A porter lifts a heavy suitcase of mass 80 kg and at the destination lowers it down by a distance of 80 cm with a constant velocity. Calculate the work done by the porter in lowering the suitcase. (Take $g = 9.8 \text{ ms}^{-2}$)
 - (a) -62720.0 J (b) +627.2 J (c) -627.2 J
 - (c) -627.2 J (d) 784.0 J
- **20.** What will be the average value of energy for a monoatomic gas in thermal equilibrium at temperature *T*?

T

T

(a)
$$k_B T$$
 (b) $\frac{3}{2} k_B$
(c) $\frac{1}{2} k_B T$ (d) $\frac{2}{3} k_B$

SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

21. The total charge enclosed in an incremental volume of 2×10^{-9} m³ located at the origin is _____ nC, if eletric flux density of its field is found as

$$D = e^{-x} \sin y \,\hat{i} - e^{-x} \cos y \,\hat{j} + 2z \,\hat{k} \, C/m^2$$

22. The position of the centre of mass of a uniform semicircular wire of radius *R* placed in *x*-*y* plane with its centre at the origin and the line joining its ends as *x*-axis

is given by $\left(0, \frac{xR}{\pi}\right)$. Then the value of |x| is _____

23. In an electric circuit, a cell of certain emf provides a potential difference of 1.25 V across a load resistance of 5 Ω . However, it provides a potential difference of 1 V across a load resistance of 2 Ω . The emf of the cell is

given by
$$\frac{x}{10}$$
 V. Then the value of x is _____.

24. In a given circuit diagram, a 5 V zener diode along with a series resistance is connected across a 50 V power supply. The minimum value of the resistance required, if the maximum zener current is 90 mA will be $___\Omega$.



25. Three students S_1 , S_2 and S_3 perform an experiment for determining the acceleration due to gravity (*g*) using a simple pendulum. They use different lengths of pendulum and record time for different number of oscillations. The observations are as shown in the table.

Student no.	Length of pendulum (cm)	No. of oscillations (n)	Total time for <i>n</i> oscillations	Time period (s)
1.	64.0	8	128.0	16.0
2.	64.0	4	64.0	16.0
3.	20.0	4	36.0	9.0

(Least count of length = 0.1 cm, least count for time = 0.1 s)

If E_1 , E_2 and E_3 are the percentage errors in 'g' for students 1, 2 and 3 respectively, then the minimum percentage error is obtained by student no. _____.

26. Three particles *P*, *Q* and *R* are moving along the vectors $\vec{A} = \hat{i} + \hat{j}$, $\vec{B} = \hat{j} + \hat{k}$ and $\vec{C} = -\hat{i} + \hat{j}$ respectively. They strike on a point and start to move in different directions. Now, particle *P* is moving normal to the plane which contains vector \vec{A} and \vec{B} . Similarly particle *Q* is moving normal to the plane which contains vector \vec{A} and \vec{B} . Similarly particle *Q* is moving normal to the plane which contains vector \vec{A} and \vec{B} . Similarly particle *Q* is moving normal to the plane which contains vector \vec{A} and \vec{C} . The angle between the

direction of motion of *P* and *Q* is $\cos^{-1}\left(\frac{1}{\sqrt{x}}\right)$. Then the value of *x* is _____.

- 27. In 5 minutes, a body cools from 75°C to 65°C at room temperature of body at the end of next 5 minutes is _____°C.
- **28.** A ray of light passing through a prism $(\mu = \sqrt{3})$ suffers minimum deviation. It is found that the angle of incidence is double the angle of refraction within the prism. Then, angle of prism is _____ (in degrees).
- **29.** The centre of a wheel rolling on a plane surface moves with a speed v_0 . A particle on the rim of the wheel at the same level as the centre will be moving at a speed $\sqrt{x} v_0$. Then the value of *x* is _____.
- 30. The area of cross-section of a railway track is 0.01 m². The temperature variation is 10°C. Coefficient of linear expansion of material of track is 10⁻⁵/°C. The energy stored per metre in the track is _____ J/m. (Young's modulus of material of track is 10¹¹ Nm⁻²)

CHEMISTRY

SECTION - A (MULTIPLE CHOICE QUESTIONS)

31. Which one of the following compounds will provide a tertiary alcohol on reaction with excess of CH₃MgBr followed by hydrolysis?





41. Match List I with List-II.

List-I

(A) Chloroprene

(B) Neoprene

(C) Acrylonitrile

(iv) $CH_2 = CH - CN$

List-II

Choose the correct answer from the options given below.

- (a) (A) (iii), (B) (iv), (C) (ii), (D) (i)
- (b) (A) (ii), (B) (i), (C) (iv), (D) (iii)
- (c) (A) (iii), (B) (i), (C) (iv), (D) (ii)
- (d) (A) (ii), (B) (iii), (C) (iv), (D) (i)
- **42.** Which purification technique is used for high boiling organic liquid compound (decomposes near its boiling point)?
 - (a) Simple distillation (b) Fractional distillation
 - (c) Reduced pressure distillation
 - (d) Steam distillation

43. Given below are the statements about diborane.

- (A) Diborane is prepared by the oxidation of NaBH₄ with I₂.
- (B) Each boron atom is in sp^2 hybridised state.
- (C) Diborane has one bridged 3 centre-2-electron bond.

(D) Diborane is a planar molecule.

- The option with correct statement(s) is
- (a) (C) only (b) (A) and (B) only
- (c) (C) and (D) only (d) (A) only
- **44.** Sulphide ion is soft base and its ores are common for metals

(A) Pb (B) Al (C) Ag (D) Mg Choose the correct answer from the options given below.

- (a) (A) and (C) only (b) (A) and (B) only
- (c) (A) and D) only (d) (C) and (D) only

45. Match List-I with List-II.

List-I	List-II
(Elements)	(Properties)
(A) Ba	(i) Organic solvent
	soluble compounds
(B) Ca	(ii) Outer electronic
	configuration $6s^2$
(C) Li	(iii) Oxalate insoluble in
	water
(D) Na	(iv) Formation of very
	strong monoacidic base

Choose the correct answer from the options given below.

- (a) (A) (i), (B)-(iv), (C)-(ii) and (D)-(iii)
- (b) (A) (ii), (B)-(iii), (C)-(i) and (D)-(iv)
- (c) (A) (iv), (B)-(i), (C)-(ii) and (D)-(iii)
- (d) (A) (iii), (B)-(ii), (C)-(iv) and (D)-(i)
- **46.** Which one of the following statements for D.I. Mendeleev, is incorrect?
 - (a) Element with atomic number 101 is named after him.
 - (b) He authored the textbook-Principles of Chemistry.
 - (c) He invented accurate barometer.
 - (d) At the time, he proposed Periodic Table of elements, structure of atom was known.
- **47.** The water having more dissolved O_2 is
 - (a) polluted water (b) water at 80°C
 - (c) boiling water (d) water at 4°C.
- **48.** When silver nitrate solution is added to potassium iodide solution then the sol produced is
 - (a) KI/NO_3^- (b) AgI/Ag^+
 - (c) $AgNO_3/NO_3^-$ (d) AgI/I^-
- **49.** Which one of the following compounds does not exhibit resonance?

(a)
$$CH_3CH_2OCH = CH_2$$
 (b) CH_2OH

- (c) $CH_3CH_2CH = CHCH_2NH_2$
- (d) CH₃CH₂CH₂CONH₂
- **50.** Isotopes(s) of hydrogen which emits low energy β^- particles with $t_{1/2}$ value > 12 years is/are
 - (a) protium (b) tritium
 - (c) deuterium (d) deuterium and tritium.

SECTION - B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

- 51. If the concentration of glucose $(C_6H_{12}O_6)$ in blood is 0.72 g L^{-1} , the molarity of glucose in blood is ______ $\times 10^{-3} \text{ M.}(\text{Nearest Integer})$ (Given : Atomic mass of C = 12, H = 1, O = 16 u)
- **52.** The total number of unpaired electrons present in $[Co(NH_3)_6]Cl_2$ and $[Co(NH_3)_6]Cl_3$ is _____.

53. A copper complex crystallising in a *ccp* lattice with a cell edge of 0.4518 nm has been revealed by employing X-ray diffraction studies. The density of a copper complex is found to be 7.62 g cm⁻³. The molar mass of copper complex is ______ g mol⁻¹. (Nearest Integer) [Given : $N_{\rm A} = 6.022 \times 10^{23} \text{ mol}^{-1}$]

54.
$$N_2O_{5(g)} \longrightarrow 2NO_{2(g)} + \frac{1}{2}O_{2(g)}$$

In the above first order reaction the initial concentration of N_2O_5 is 2.40×10^{-2} mol L⁻¹ at 318 K. The concentration of N_2O_5 after 1 hour was 1.60×10^{-2} mol L⁻¹. The rate constant of the reaction at 318 K is ______ $\times 10^{-3}$ min⁻¹. (Nearest Integer) [Given : log3 = 0.477, log5 = 0.699]

- 55. Methylation of 10 g of benzene gave 9.2 g of toluene. Calculate the percentage yield of toluene ______.(Nearest Integer)
- 56. Assume a cell with the following reaction $Cu_{(s)} + 2Ag^+(1 \times 10^{-3} \text{ M}) \rightarrow Cu^{2+}(0.250 \text{ M}) + 2Ag_{(s)}$ $E_{cell}^\circ = 2.97 \text{ V}$ E_{cell} for the above reaction is _____ V. (Nearest Integer) [Given : log 2.5 = 0.3979, T = 298 K]

[Given : log 2.5 = 0.5979, 1 = 298 K]

- 57. If the standard molar enthalpy change for combustion of graphite powder is -2.48×10^2 kJ mol⁻¹, the amount of heat generated on combustion of 1 g of graphite powder is _____ kJ. (Nearest Integer)
- **58.** Number of electrons that Vanadium (Z = 23) has in *p*-orbitals is equal to _____.
- **59.** The number of acyclic structural isomers (including, geometrical isomers) for pentene are _____.
- **60.** Value of K_p for the equilibrium reaction $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ at 288 K is 47.9. The K_c for this reaction at same temperature is ______. (Nearest Integer) (R = 0.083 L bar K⁻¹ mol⁻¹)

MATHEMATICS

SECTION - A (MULTIPLE CHOICE QUESTIONS)

- **61.** Let $A = [a_{ij}]$ be a real matrix of order 3×3 , such that $a_{i1} + a_{i2} + a_{i3} = 1$, for i = 1, 2, 3. Then, the sum of all the entries of the matrix A^3 is equal to (a) 1 (b) 9 (c) 3 (d) 2
- **62.** Four dice are thrown simultaneously and the numbers shown on these dice are recorded in 2×2 matrices. The probability that such formed matrices have all different entries and are non-singular, is

(a)
$$\frac{45}{162}$$
 (b) $\frac{23}{81}$ (c) $\frac{43}{162}$ (d) $\frac{22}{81}$

63. The number of solutions of $\sin^7 x + \cos^7 x = 1$, $x \in [0, 4\pi]$ is equal to (a) 9 (b) 5 (c) 11 (d) 7

- **64.** Let three vectors \vec{a}, \vec{b} and \vec{c} be such that $\vec{a} \times \vec{b} = \vec{c}$, $\vec{b} \times \vec{c} = \vec{a}$ and $|\vec{a}| = 2$. Then, which one of the following is not true?
 - (a) $[\vec{a} \, \vec{b} \, \vec{c}] + [\vec{c} \, \vec{a} \, \vec{b}] = 8$
 - (b) Projection of \vec{a} on $(\vec{b} \times \vec{c})$ is 2.
 - (c) $|3\vec{a} + \vec{b} 2\vec{c}|^2 = 51$
 - (d) $\vec{a} \times ((\vec{b} + \vec{c}) \times (\vec{b} \vec{c})) = \vec{0}$
- 65. Let *n* denote the number of solutions of the equation $z^2 + 3\overline{z} = 0$, where *z* is a complex number. Then the value of $\sum_{k=0}^{\infty} \frac{1}{n^k}$ is equal to (a) $\frac{3}{2}$ (b) 2 (c) 1 (d) $\frac{4}{3}$
- 66. Let L be the line of intersection of planes $\vec{r} \cdot (\hat{i} \hat{j} + 2\hat{k}) = 2$ and $\vec{r} \cdot (2\hat{i} + \hat{j} \hat{k}) = 2$. If $P(\alpha, \beta, \gamma)$ is the foot of perpendicular on L from the point (1, 2, 0), then the value of $35(\alpha + \beta + \gamma)$ is equal to (a) 134 (b) 101 (c) 143 (d) 119
- **67.** Which of the following Boolean expressions is not a tautology?
 - (a) $(p \Rightarrow \neg q) \lor (\neg q \Rightarrow p)$ (b) $(p \Rightarrow q) \lor (\neg q \Rightarrow p)$ (c) $(q \Rightarrow p) \lor (\neg q \Rightarrow p)$ (d) $(\neg p \Rightarrow q) \lor (\neg q \Rightarrow p)$

68. Let [x] denote the greatest integer less than or equal to x. Then, the values of x ∈ R satisfying the equation [e^x]² + [e^x + 1] - 3 = 0 lie in the interval
(a) [0, 1/e)
(b) [0, log_e2)

(c)
$$[\log_e 2, \log_e 3)$$
 (d) $[1, e)$

69. Let $E_1: \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, a > b. Let E_2 be another ellipse such that it touches the end points of major axis of E_1 and the foci of E_2 are the end points of minor axis of E_1 . If E_1 and E_2 have same eccentricities, then its value is

(a)
$$\frac{-1+\sqrt{6}}{2}$$
 (b) $\frac{-1+\sqrt{2}}{2}$
(c) $\frac{-1+\sqrt{8}}{2}$ (d) $\frac{-1+\sqrt{2}}{2}$

70. If the domain of the function $f(x) = \frac{\cos^{-1}\sqrt{x^2 - x + 1}}{\sqrt{\sin^{-1}\left(\frac{2x - 1}{2}\right)}}$

 $\sqrt{5}$

is the interval $(\alpha, \beta]$, then $\alpha + \beta$ is equal to

(a) 1 (b)
$$\frac{1}{2}$$
 (c) 2 (d) $\frac{3}{2}$

71. Let $f: R \to R$ be defined as

$$f(x) = \begin{cases} \frac{x^3}{(1 - \cos 2x)^2} \log_e \left(\frac{1 + 2xe^{-2x}}{(1 - xe^{-x})^2} \right), & x \neq 0 \\ \alpha, & x = 0 \end{cases}$$

If *f* is continuous at x = 0, then α is equal to (a) 2 (b) 0 (c) 3 (d) 1





72. If
$$\int_{0}^{100\pi} \frac{\sin^2 x}{e^{\left(\frac{x}{\pi} - \left[\frac{x}{\pi}\right]\right)}} dx = \frac{\alpha \pi^3}{1 + 4\pi^2}, \alpha \in R$$
, where [x] is the

greatest integer less than or equal to x, then the value of α is

- (a) $200(1 e^{-1})$ (b) $150(e^{-1} 1)$ (c) 50(e - 1) (d) 100(1 - e)
- 73. Let a line L: 2x + y = k, k > 0 be a tangent to the hyperbola $x^2 y^2 = 3$. If *L* is also a tangent to the parabola $y^2 = \alpha x$, then α is equal to (a) -12 (b) 24 (c) 12 (d) -24
- 74. Let the circle $S: 36x^2 + 36y^2 108x + 120y + C = 0$ be such that it neither intersects nor touches the co-ordinate axes. If the point of intersection of the lines, x - 2y = 4 and 2x - y = 5 lies inside the circle *S*, then
 - (a) 81 < C < 156 (b) 100 < C < 16525 13

(c)
$$100 < C < 156$$
 (d) $\frac{25}{9} < C < \frac{15}{3}$

- 75. If the shortest distance between the straight lines 3(x-1)=6(y-2)=2(z-1) and $4(x-2)=2(y-\lambda)=(z-3)$, $\lambda \in R$ is $\frac{1}{\sqrt{38}}$, then the integral value of λ is equal to
 - (a) 3 (b) 5 (c) 2 (d) -1
- 76. Let a vector \vec{a} be coplanar with vectors $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - \hat{j} + \hat{k}$. If \vec{a} is perpendicular to $\vec{d} = 3\hat{i} + 2\hat{j} + 6\hat{k}$, and $|\vec{a}| = \sqrt{10}$. Then, a possible value of $[\vec{a} \ \vec{b} \ \vec{c}] + [\vec{a} \ \vec{b} \ \vec{d}] + [\vec{a} \ \vec{c} \ \vec{d}]$ is equal to (a) -38 (b) -29 (c) -40 (d) -42
- 77. Let y = y(x) be the solution of the differential equation $\csc^2 x dy + 2 dx = (1 + y \cos 2x) \csc^2 x dx$, with

$$y\left(\frac{\pi}{4}\right) = 0$$
. Then, the value of $(y(0) + 1)^2$ is equal to
(a) $e^{-\frac{1}{2}}$ (b) $e^{-\frac{1}{2}}$ (c) $e^{\frac{1}{2}}$ (d) e^{-1}

(a)
$$e^{-2}$$
 (b) e (c) e^{2} (c)

78. Let $f: R \to R$ be defined as

$$f(x) = \begin{cases} -\frac{4}{3}x^3 + 2x^2 + 3x, & x > 0\\ 3xe^x, & x \le 0 \end{cases}$$

Then f is increasing function in the interval

(a)
$$\left(-1, \frac{3}{2}\right)$$
 (b) $\left(-\frac{1}{2}, 2\right)$
(c) $(0, 2)$ (d) $(-3, -1)$

- **79.** The values of λ and μ such that the system of equations x + y + z = 6, 3x + 5y + 5z = 26, $x + 2y + \lambda z = \mu$ has no solution, are
 - (a) $\lambda = 2, \mu \neq 10$ (b) $\lambda \neq 2, \mu = 10$
 - (c) $\lambda = 3, \mu = 5$ (d) $\lambda = 3, \mu \neq 10$
- **80.** Let S_n denote the sum of first *n*-terms of an arithmetic progression. If $S_{10} = 530$, $S_5 = 140$, then $S_{20} S_6$ is equal to (a) 1862 (b) 1852 (c) 1842 (d) 1872

SECTION - B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

- **81.** The sum of all the elements in the set $\{n \in \{1, 2, \dots, 100\} | \text{ H.C.F. of } n \text{ and } 2040 \text{ is } 1\}$ is equal to _____.
- **82.** Let $f: R \to R$ be a function defined as

$$f(x) = \begin{cases} 3\left(1 - \frac{|x|}{2}\right), & \text{if } |x| \le 2\\ 0, & \text{if } |x| > 2 \end{cases}$$

Let $g : R \to R$ be given by g(x) = f(x + 2) - f(x - 2). If n and m denote the number of points in R, where g is not continuous and not differentiable, respectively, then n + m is equal to _____.

- **83.** The area (in sq. units) of the region bounded by the curves $x^2 + 2y 1 = 0$, $y^2 + 4x 4 = 0$ and $y^2 4x 4 = 0$, in the upper half plane is _____.
- 84. If the constant term, in binomial expansion of $\left(2x^r + \frac{1}{x^2}\right)^{10}$ is 180, then *r* is equal to _____.
- **85.** Let *A* = {0, 1, 2, 3, 4, 5, 6, 7}. Then the number of bijective functions *f* : *A* → *A* such that f(1) + f(2) = 3 f(3) is equal to _____.
- **86.** Let y = y(x) be the solution of the differential equation

$$\left((x+2)e^{\left(\frac{y+1}{x+2}\right)} + (y+1)\right)dx = (x+2)dy, y(1) = 1.$$

If the domain of y = y(x) is an open interval (α, β) , then $|\alpha + \beta|$ is equal to _____.

- **87.** If the digits are not allowed to repeat in any number formed by using the digits 0, 2, 4, 6, 8, then the number of all numbers greater than 10,000 is equal to _____.
- **88.** The number of elements in the set $\{n \in \{1, 2, 3, \dots, 100\} \mid (11)^n > (10)^n + (9)^n\}$ is _____
- **89.** Consider the following frequency distribution:

Class	0-6	6-12	12-18	18-24	24-30
Frequency	а	Ь	12	9	5

If mean $=\frac{309}{22}$ and median = 14, then the value $(a - b)^2$ is equal to _____.

s equal to _____

90. Let
$$A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
. Then the number of 3 × 3 matrices

B with entries from the set $\{1, 2, 3, 4, 5\}$ and satisfying AB = BA is _____.



HINTS & EXPLANATIONS

1. (**b**): (A) If $\omega L > \frac{1}{\omega C}$, means $X_L > X_C$, ϕ is positive, so current lags behind the applied emf.

(B) If $\omega L = \frac{1}{\omega C}$, $X_L = X_C$, so current is in phase with emf.

(C) If $\omega L < \frac{1}{\omega C}$, $X_L < X_C$, ϕ is negative, so current leads

the emf.

(D) At resonant frequency, maximum current occurs.
 (A)→(ii), (B)→(i), (C)→(iv), (D)→(iii)

2. (c) : In ferromagnetic material, all the molecular magnetic dipoles are pointed in the same direction. When the ferromagnetic material heated, all magnetic dipoles are distributed and get disoriented. Due to which the net magnetic dipole moment becomes very less, so that they behave as paramagnetic material. So, statement I is true.

3. (d) : The band gap energy is

$$\Delta E = \frac{hc}{\lambda_0}$$

where, *h* is Planck's constant rod, *c* is speed of light and λ_0 is the threshold wavelength.

$$\therefore \quad \Delta E = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{621 \times 10^{-9}} \text{ J}$$
$$\Delta E = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{621 \times 10^{-9} \times 1.6 \times 10^{-19}} = 2 \text{ eV}$$

4. (b) : Length of Copper rod, $L_1 = 25$ cm, $\rho_{Cu} = 1.7 \times 10^{-8} \Omega$ m Area of copper rod, $A_1 = 3$ mm²,

Length of Aluminium rod, $L_2 = 25$ cm, $\rho_{Al} = 2.6 \times 10^{-8}$ W m Area of Aluminium rod = 3 mm²

Here both the wires are connected in parallel as shown in figure.

$$A - \underbrace{\begin{array}{c} Cu \\ Al \end{array}}_{A} - \underbrace{\begin{array}{c} Cu \\ Al \end{array}}_{B}$$

$$R_{1} = \rho_{1} \frac{L_{1}}{A_{1}} \text{ and } R_{2} = \rho_{2} \frac{L_{2}}{A_{2}}$$

$$\frac{1}{R_{eq}} = \frac{1}{R_{1}} + \frac{1}{R_{2}} = \frac{A_{1}}{\rho_{1}L_{1}} + \frac{A_{2}}{\rho_{2}L_{2}}$$

$$\frac{1}{R_{eq}} = \frac{A}{L} \left(\frac{1}{\rho_{1}} + \frac{1}{\rho_{2}}\right) = \frac{3 \times 10^{-6}}{0.25} \left(\frac{10^{8}}{1.7} + \frac{10^{8}}{2.6}\right)$$

$$\frac{1}{R_{eq}} = \frac{3 \times 10^{-6} \times 10^{8}}{0.25} \left[\frac{2.6 + 1.7}{2.6 \times 1.7}\right] = 1167.42$$
or $R_{eq} = 0.858 \times 10^{-3} \Omega$

$$\therefore R_{eq} = 0.858 \text{ m}\Omega$$

5. (b) : In case of capacitor the current lags by $\pi/2$ by emf. So, option (b) is correct.

6. (a) : Given,
$$\vec{A} = \hat{i} + \hat{j} + \hat{k}$$
, $\vec{B} = \hat{i} + \hat{j}$
Projection of vector A over vector $B = (A \cdot B)B$
 $= (\hat{i} + \hat{j} + \hat{k}) \cdot (\hat{i} + \hat{j}) \frac{[\hat{i} + \hat{j}]}{\sqrt{2}}$
 $= \frac{(1+1)(\hat{i} + \hat{j})}{\sqrt{2}}$ (: $\hat{i} \cdot \hat{i} = 1, \hat{j} \cdot \hat{j} = 1$)
 $= \sqrt{2}(\hat{i} + \hat{j})$
7. (c) : Moment of inertia of ring $= mr^2$
Moment of inertia of solid cylinder $= \frac{1}{2}mr^2$
Moment of inertia of solid sphere $= \frac{2}{5}mr^2$
The acceleration is given by, $a = \frac{g\sin\theta}{1 + \frac{K^2}{r^2}}$...(i)
For ring, $K = r$
For solid cylinder, $K = \frac{r}{\sqrt{2}}$
For solid sphere, $K = \sqrt{\frac{2}{5}}r$
Now, using eq. (i),
 $a_{ring} = \frac{g\sin\theta}{2} = 0.5 g\sin\theta$
 $a_{solid cylinder} = \frac{g\sin\theta}{1 + \frac{1}{2}} = \frac{2}{3}g\sin\theta = 0.66g\sin\theta$

$$a_{\text{solid sphere}} = \frac{g\sin\theta}{1+\frac{2}{5}} = \frac{5}{7}g\sin\theta = 0.7g\sin\theta$$

So, the sphere has the greatest and ring has least velocity of centre of mass at the bottom of inclined plane.

8. (c) : Radius coverage, $R = 150 \text{ km} = 150 \times 1000 \text{ m}$ Population density = 2000/km² Radius of earth, $R_e = 6.5 \times 10^6 \text{ m}$ Let the height is H. So, radius $R = \sqrt{2R_eH}$ $150 \times 1000 = \sqrt{2 \times 6.5 \times 10^6 \times H}$ $(1000 \times 150)^2 = 2 \times 6.5 \times 10^6 H$ H = 1731 m.

Population covered = $\pi R^2 \times$ population density = $3.14(150)^2 \times 2000 = 1413 \times 10^5$



9. (b) : Intensity of sunlight, $I = 0.092 \text{ W/m}^2$ $\varepsilon_0 = 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$ Intensity is given by

$$I = \frac{1}{2} \varepsilon_0 E_0^2 = \frac{1}{2} \frac{B_0^2 c}{\mu_0}$$

$$\frac{1}{\sqrt{\mu_0 \varepsilon_0}} = c \Longrightarrow \frac{1}{\mu_0} = \varepsilon_0 c^2$$

So, $I = \frac{1}{2} B_0^2 c \varepsilon_0 c^2 = \frac{1}{2} B_0^2 \varepsilon_0 c^3$
 $B_0 = \sqrt{\frac{2I}{\varepsilon_0 c^3}} = \sqrt{\frac{2 \times 0.092}{8.85 \times 10^{-12} \times (3 \times 10^8)^3}}$
 $B_0 = 2.77 \times 10^{-8} \text{ T}$

10. (a) : Linear charge density, $\lambda = 3 \times 10^{-6}$ C/m Net force, F = 4 N r = 10 mm, x = 12 mm

Let the charge on dipole is q

$$F_{1} = \frac{2k\lambda}{r} \cdot q$$

$$F_{2} = \frac{2k\lambda q}{r+x}$$

$$F_{net} = F_{1} - F_{2} = \frac{2K\lambda q}{r} - \frac{2K\lambda q}{r+x}$$

$$F_{net} = \frac{2K\lambda q \cdot x}{r(r+x)}$$

$$4 = \frac{2 \times 9 \times 10^{9} \times 3 \times 10^{-6} \times 2 \times 10^{-3} \times q}{10 \times 10^{-3} \times 12 \times 10^{-3}}$$

 $q = 4.44 \times 10^{-6} \text{ C} = 4.44 \ \mu\text{C}$

11. (a) : Mass number, A = 184 $Q_{\text{value}} = 5.5 \text{ MeV}$ Let the velocity of α -particle is *v* and for 180 *m*, it is *v'*.

$$\bigcap_{\text{Rest}}^{184 m} \qquad \bigvee^{\prime} \stackrel{180 m}{\longrightarrow} \qquad \stackrel{4 m}{\longrightarrow} \qquad \stackrel{\nu}{\longrightarrow} \qquad \qquad$$

Use conservation of momentum, 184 $m \times 0 = 180 mv' - 4mv$

$$\nu' = \frac{4\nu}{180} \qquad \dots (i)$$

Now using conservation of energy,

$$\frac{1}{2}(4m)v^{2} + \frac{1}{2}(180m)v'^{2} = 5.5 \text{ MeV}$$
$$\frac{1}{2} \cdot 4mv^{2} \left[1 + 45 \times \left(\frac{4}{180}\right)^{2} \right] = 5.5 \text{ MeV} \text{ (Using (i))}$$
Here $\text{K} \cdot \text{E}_{\alpha} = \frac{1}{2}(4mv^{2})$ $\text{K} \cdot \text{E}_{\alpha} \left(1 + 45 \times \left(\frac{4}{180}\right)^{2} \right) = 5.5 \text{ MeV}$

$$K \cdot E_{\alpha} = \frac{5.5}{1 + 45 \left(\frac{4}{180}\right)^2} = 5.38 \text{ MeV}$$

12. (d) : The minimum velocity of body is escape velocity as it goes to infinity $(A^{2}y_{2})$

$$v_1 = \sqrt{\frac{2GM}{R_e}} \qquad \dots (i)$$

Let it reaches to distance r with velocity v_2 and time t. Use conservation of energy

$$\frac{1}{2}m\left(\frac{2GM}{R_e}\right) - \frac{GMm}{R_e} = \frac{1}{2}mv_2^2 - \frac{GMm}{R+r}$$

$$\frac{1}{2}mv_2^2 = \frac{GMm}{R+r}$$

$$v_2 = \sqrt{\frac{2GM}{R+r}}$$
Velocity is given by
$$v = \frac{dr}{dt}$$

$$\sqrt{2GM} \int_0^t dt = \int_{R_e}^{R_e+h} \sqrt{R+r} dr$$

$$\sqrt{2GM} \times t = \frac{2}{3} \left[(R+r)^{3/2} \right]_{R_e}^{R_e+h}$$

$$t = \frac{2}{3} \sqrt{\frac{R_e^3}{2GM}} \left[\left(1 + \frac{h}{R_e} \right)^{3/2} - 1 \right]$$

$$\therefore \quad t = \frac{1}{3}\sqrt{\frac{2R_e}{g}} \left[\left(1 + \frac{h}{R_e}\right)^{3/2} - 1 \right]$$

$$\tan \phi = \frac{B_V}{B_H} \qquad \dots(i)$$

and
$$\tan \phi' = \frac{B_V}{B_H \cos \alpha} = \frac{\tan \phi}{\cos \alpha} \qquad (From (i))$$

True dip circle as shown in figure (i).
$$B_V \qquad Figure (i)$$

$$Figure (i)$$

Apparent dip circle as shown in figure (ii).

So, $\tan \phi = \tan \phi' \cos \alpha$ (0 < cos α \therefore $\tan \phi < \tan \phi'$ Hence, $\phi < \phi'$

 $\left(\because g = \frac{GM}{R_e^2} \right)$

13. (b) : Let the apparent dip is at α with true dip B_{V}

 \dot{B}_V Figure (0 < cos α < 1)

Figure (ii)

🝸 unacademy

...(i)

...(ii)

14. (c) : According to question, $r + r' = 90^{\circ}$ and $r + r' + 90^{\circ} = 180^{\circ}$ $r' = 90^{\circ} - r = 90^{\circ} - i$ By Snell's law $n_1 \sin i = n_2 \sin r'$, From eq. (i), we have $n_1 \sin i = n_2 \sin(90^{\circ} - i)$ $n_1 \sin i = n_2 \cos i \implies \tan i = \frac{n_2}{2}$

 $n_1 \sin i = n_2 \cos i \implies \tan i = \frac{n_2}{n_1}$ Also $\sin C = \frac{n_2}{n_1}$ where, C is critical angle.

From eq. (ii), we get $\sin C = \tan i$ $C = \sin^{-1}(\tan i) = \sin^{-1}(\tan r)$

15. (b) : Initial time period of a simple pendulum, $T_1 = T_0$, and length = *L*

Final length, $L' = \frac{L}{16}$

Let the new time period is T'. The time period of simple pendulum is given by

 $T = 2\pi \sqrt{\frac{L}{g}}$

So, $T \propto \sqrt{L}$

$$\therefore \quad \frac{T'}{T_0} = \sqrt{\frac{L}{16L}} = \frac{1}{4} \implies T' = \frac{T_0}{4}$$

16. (c) : Mass of bullet, m = 4 g Mass of gun, M = 4 kg Muzzle speed of bullet, v = 50 m/s Let the recoil velocity of gun is *V*. Use conservation of momentum

mv = MV $0.004 \times 50 = 4 V$ V = 0.05 m/sAlso, Impulse = Change in momentum MV - mv $= MV - 0 = 4 \times 0.05 = 0.2 \text{ kg m/s}$

17. (d): Given,
$$x(t) = A\sin\omega t + B\cos\omega t$$

As,
$$v = \frac{dx}{dt} = A\omega\cos\omega t - B\omega\sin\omega t$$

At
$$t = 0$$
, $x = B$, and $v = A\omega$

From the graph, $A_{\text{net}} = \sqrt{A^2 + B^2}$

or $\tan \alpha = \frac{B}{A} \Longrightarrow \alpha = \tan^{-1} \left(\frac{B}{A} \right)$

$$x(t) = \sqrt{A^2 + B^2} \left[\frac{A}{\sqrt{A^2 + B^2}} \sin \omega t + \frac{B}{\sqrt{A^2 + B^2}} \cos \omega t \right]$$

Let
$$\frac{A}{\sqrt{A^2 + B^2}} = \sin\phi$$
; $\frac{B}{\sqrt{A^2 + B^2}} = \cos\phi$
 $\Rightarrow x(t) = C[\sin\phi\sin\omega t + \cos f \cos\omega t]$
 $\Rightarrow x(t) = C\cos(\omega t - \phi)$
As, $C = \sqrt{A^2 + B^2} = \sqrt{\frac{[v(0)]^2}{\omega^2} + x(0)^2}$
 $\phi = 90 - \alpha$,
 $\tan\phi = \cot\alpha = \frac{A}{B} = \frac{v(0)}{x(0)\omega}$
 $\phi = \tan^{-1}\left[\frac{V(0)}{\omega x(0)}\right]$

18. (d): The de Broglie wavelength is

$$\lambda = \frac{h}{\sqrt{2mqV}}$$

where, h is Planck's constant, m is mass and q is charge, V is potential difference.

So,
$$\lambda \propto \frac{1}{\sqrt{2mV}}$$

As potential difference is same, so

$$\therefore \frac{\lambda_e}{\lambda_p} = \sqrt{\frac{m_p}{m_e}}$$

19. (c) : Mass of suitcase, m = 80 kgDistance = h = 80 cm, $g = 9.8 \text{ m/s}^2$ Let the work done by porter is W_{porter} . According to the work energy theorem $W_{\text{porter}} + W_{mg} = \Delta KE$ $W_{\text{porter}} = -W_{mg} = -mgh$ ($\because \Delta KE = 0$) = $(80 \times 9.8 \times 0.8)$ = -627.2 J

20. (b) : The average value of energy for a monoatomic gas in thermal equilibrium at temperature *T* is

$$U = \frac{3}{2} k_B T$$

(:: v = 0)

where, k_B is Boltzmann constant.

21. (4) : The electric flux density is

$$\vec{D} = \frac{\text{Charge}}{\text{Area}}\hat{r} = \frac{Q}{4\pi r^2}\hat{r} = \varepsilon_0 \left[\frac{Q}{4\pi\varepsilon_0 r^2}\right]\hat{r}$$

Given,
$$D = e^{-x} \sin y \hat{i} - e^{-x} \cos y \hat{j} + 2z\hat{k} C/m^2$$

As,
$$E = \frac{\vec{D}}{\varepsilon_0} = \frac{e^{-x} \sin y \,\hat{i} - e^{-x} \cos y \,\hat{j} + 2z \,\hat{k}}{\varepsilon_0}$$

According to Gauss's theorem

$$\frac{\rho}{\varepsilon_0} = \left(\frac{\partial}{\partial x}\hat{i} + \frac{\partial}{\partial y}\hat{j} + \frac{\partial}{\partial z}\hat{k}\right)\cdot\vec{E}$$



or
$$\frac{\rho}{\varepsilon_0} = \left(\frac{\partial}{\partial x}\hat{i} + \frac{\partial}{\partial y}\hat{j} + \frac{\partial}{\partial z}\hat{k}\right)\cdot\frac{\vec{D}}{\varepsilon_0}$$

Also, $\rho = \frac{\partial}{\partial x}(e^{-x}\sin y) + \frac{\partial}{\partial y}(-e^{-x}\cos y) + \frac{\partial}{\partial z}(2z)$
or $\rho = -e^{-x}\sin y + e^{-x}\sin y + 2$
At origin $\rho = 2 \text{ C/m}^3$
The total charge enclosed, $q = \rho \times \text{Volume} = 2 \times 2 \times 10^{-9}$
 $= 4 \times 10^{-9} \text{ C}$
 $\therefore q = 4 \text{ nC}$

22. (2): Let M is the mass, R is the radius and O is the centre.

 $\frac{\text{Mass}}{\text{Length}} = \frac{M}{\pi R}$

Consider an elementary portion, so mass of elementary



The coordinates of centre of mass are

$$X = \frac{1}{M} \int x \, dm = \frac{1}{M} \int_{0}^{\pi} (R \cos \theta) \left(\frac{M}{\pi}\right) d\theta = \frac{R}{\pi} [\sin \theta]_{0}^{\pi}$$

So, $X = 0$

Similarly,
$$Y = \frac{1}{M} \int y \, dm = \frac{1}{M} \int_{0}^{\pi} R \sin \theta \frac{M}{\pi} \cdot d\theta = \frac{2R}{\pi}$$

On comparing the given value with $\frac{xR}{\pi}$, we get, x = 2.

π

23. (15) : Let the emf is *E* and internal resistance is *r*.
Case I : Current,
$$I = \frac{V}{r} = \frac{1.25}{5}$$

As, $\varepsilon - Ir = V$
 $\varepsilon - \frac{1.25}{5}r = 1.25$...(i)

Case II : Current, $I = \frac{V}{r} = \frac{1}{2}$ Also, $\varepsilon - Ir = V$ or $\varepsilon - \frac{1}{2}r = 1$...(ii)

By solving eqns. (i) and (ii), we get $\varepsilon = 1.5 \text{ V}, r = 1 \Omega$

On comparing the given value, with $\frac{x}{10}$ V, we get x = 15

$$\left(:: \frac{15}{10} \text{V or } 1.5 \text{V}\right) \qquad t = \frac{1}{K} \ln \left(\frac{\theta_2}{\theta_1}\right)$$

24. (500) : Given,
$$I_1 = 90$$
 mA, $I_2 = \frac{5}{R_L}$...(i)

Voltage across *R*, V' = 50 - 5 = 45 V

So,
$$R = \frac{45}{I_1 + I_2}$$

 $R = \frac{45}{0.09 + \frac{5}{R_L}}$ (Using (i))

The current in zener is maximum when $R_L = \infty$ So, $I_2 = 0$, $I_1 = I$

$$\therefore \quad R = \frac{45}{0.09} = 500 \,\Omega$$

25. (1): The time period of simple pendulum is given by

$$T = 2\pi \sqrt{\frac{L}{g}}$$

where, L is length of pendulum and g is the acceleration due to gravity.

Also,
$$T^2 = 4\pi^2 \frac{L}{g}$$

 $g = 4\pi^2 \frac{L}{T^2}$

Percentage error, $\frac{\Delta g}{g} \times 100 = \frac{\Delta L}{L} \times 100 + \frac{2\Delta T}{nT} \times 100$

 ΔL and ΔT is same for all so, $\frac{\Delta g}{\tilde{L}}$ is minimum for highest value of *L*, *n* and *T*. So, the minimum percentage error is obtained by student 1.

26. (3): Given,
$$\vec{A} = \hat{i} + \hat{j}$$
, $\vec{B} = \hat{j} + \hat{k}$, $\vec{C} = -\hat{i} + \hat{j}$

$$\check{n}_1 = \frac{\vec{A} \times \vec{B}}{|\vec{A} \times \vec{B}|} = \frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$$
$$\hat{n}_2 = \frac{\vec{A} \times \vec{C}}{|\vec{A} \times \vec{C}|} = \frac{2\hat{k}}{2} = \hat{k}$$

Angle between \hat{n}_1 and \hat{n}_2 is

$$\cos\theta = \frac{\hat{n}_1 \cdot \hat{n}_2}{|\hat{n}_1||\hat{n}_2|} = \frac{\frac{1}{\sqrt{3}}}{(1)(1)} = \frac{1}{\sqrt{3}} \text{ or } \theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

On comparing the given value with $\cos^{-1}\left(\frac{1}{\sqrt{r}}\right)$, We get, x = 3.

27. (57) : In 5 min, a body cools from temperature 75°C to 65°C.

Room temperature = $25^{\circ}C$ By Newton's law of cooling,

$$t = \frac{1}{K} \ln \left(\frac{\theta_2 - \theta_n}{\theta_1 - \theta_n} \right)$$

 $\mu = \sqrt{3}$

...(i)

where, t is time, θ_2 is final temperature, θ_1 is initial temperature, θ_n = temperature of surroundings.

So
$$5 = \frac{1}{K} \ln \left(\frac{65 - 25}{75 - 25} \right)$$
 ...(i)

Let the temperature is θ' after 5 minutes.

Now,
$$5 = \frac{1}{K} \ln \left(\frac{\theta' - 25}{65 - 25} \right)$$
 ...(ii)

From eq. (i) and (ii), we have

$$\ln\left(\frac{40}{50}\right) = \ln\left(\frac{\theta' - 25}{40}\right)$$
$$\frac{4}{5} = \frac{\theta' - 25}{40}$$
or $\theta' = 57^{\circ}C$

28. (60) : Given, with in the prism, i = 2rAccording to Snell's law,

$$\mu = \frac{\sin i}{\sin r} = \frac{\sin 2r}{\sin r}$$

. .

 $\sqrt{3} = \frac{2\sin r \cos r}{\sin r}$

$$\cos r = \frac{\sqrt{3}}{2} \implies r = 30^{\circ}$$

For minimum deviation, angle of prism

 $A = 2r = 2 \times 30^{\circ} = 60^{\circ}$

29. (2) : Speed at centre of a rolling wheel = v_0 $v_0 = R\omega$

$$\sim$$

At point *P*,
$$v = R\omega = \frac{v_0}{\omega} \cdot \omega = v_0$$
 (Using (i))

Resultant speed,
$$v = \sqrt{v_0^2 + v_0^2} = \sqrt{2}v_0$$

On comparing given value with $\sqrt{x}v_0$, we get x = 2.

30. (5) : Given,
$$A = 0.01 \text{ m}^2$$
, $\alpha = 10^{-5} \text{ °C}$, $Y = 10^{11} \text{ N/m}^2$

As, Energy = $\frac{Y}{2} \times (\text{Strain})^2 \times \text{Area} \times \text{length}$ $\frac{\text{Energy}}{\text{Length}} = \frac{Y}{2} \times \left(\frac{\Delta l}{l}\right)^2 \times A = \frac{Y}{2}A \times (\alpha \cdot \Delta T)^2$

$$= \frac{10^{11}}{2} \times 0.01(10^{-5} \times 10)^2 = 5 \text{ J/m} \qquad (\because \Delta l = l\alpha \,\Delta T)$$

The energy stored per meter in the track is 5 J/m.



33. (b) : AlCl₃ is used as catalyst in Friedel-Crafts reaction which is acidic in nature and aniline gives lone pair to AlCl₃ and converts to cation.



Thus, benzene ring is deactivated towards Friedel-Crafts reaction. Hence, aniline does not undergo Friedel-Crafts reaction.

34. (a) : Thiamine is known as vitamin B_1 while pyridoxine is vitamin B₆.

35. (c) :
$$\Delta T_f \propto i$$

So, greater the value of *i*, greater will depression in freezing point and lower will be freezing point.

Solution	<i>(i)</i>
KI	2
K_2SO_4	3
$Al_2(SO_4)_3$	5
$C_{6}H_{12}O_{6}$	1(No ionization)



36. (**b**) : Group-15 hydrides: NH₃, PH₃, AsH₃, SbH₃, BiH₃

As we move down the group, reducing power increases.



This set of ions is both paramagnetic and coloured.

39. (a) :
$$CH_3 - CH_2 - C = CH - CH_2 - CH_3$$

 $\downarrow CH_2 - CH_3$
 $_3$ -Ethvlhex-3-ene

The compound neither shows geometrical (because both alkyl groups are same on one side of the carbon attached to double bond) nor optical isomerism (no asymmetric carbon is present).



42. (c) : The compounds which have high boiling point and generally decompose before their boiling point is reached, can be purified by distillation under reduced pressure which lowers the boiling point of most liquids. So, these can be collected before decomposition.

43. (d) : The boron atom is sp^3 -hybridised in diborane. Diborane is not planar in structure as 2 hydrogen atoms are present above and below the plane containing four terminal bonds between B and H. Two 3-centre-2-electron bonds or banana bonds are present in diborane.

 $2NaBH_4 + I_2 \longrightarrow B_2H_6 + 2NaI + H_2$

- **44.** (a) : Sulphide ores \Rightarrow PbS, Ag₂S, CuFeS₂, ZnS.
- **45.** (b) : Ba $[Xe]6s^2$
 - Ca CaC_2O_4 is highly insoluble in water.
 - Li Organic solvent soluble compounds due to covalent nature
 - Na Formation of very strong monoacidic base *eg.*, NaOH

46. (d) : Preliminary work for his great textbook "Principles of Chemistry" led Mendeleev to propose the periodic law and to construct his Periodic Table of elements. At that time, the structure of atom was unknown and Mendeleev's

idea to consider that the properties of the elements were in someway related to their atomic masses was a very imaginative one.

The element with atomic number 101 is named as Mendelevium.

47. (d): Solubility of oxygen increases with decrease in temperature.

48. (d) : $AgNO_3 + KI \longrightarrow AgI/I^-$

49. (c) : NH_2 has no conjugation between π -bond and lone-pair of N. Hence, there will be no resonance in this compound. Rest all will show resonance.

50. (b) : Only tritium is radioactive and emits low energy β particles ($t_{1/2} = 12.33$ years)

51. (4):
$$M = \frac{W_{\text{solute}}}{M_{\text{solute}} \times V_{\text{soln}}(\text{in L})} = \frac{0.72}{180}$$

= 0.004 = 4 × 10⁻³ M

52. (1) : Complex

(i)
$$[Co(NH_3)_6]Cl_2 \Rightarrow Co^{2+} = 3d^7 (t_{2g}^6 e_g^1),$$

unpaired electron = 1

(ii)
$$[Co(NH_3)_6]Cl_3 \Rightarrow Co^{3+} = 3d^6 (t_{2g}^6 e_g^0),$$

unpaired electron = 0

Total unpaired electrons =1

53. (106) :
$$d = \left\{ \frac{Z \times M}{N_A \times \text{Volume}} \right\}$$

$$7.62 = \frac{4 \times M}{6.022 \times 10^{23} \times [0.4518 \times 10^{-7}]^3}$$
$$M = \frac{7.62 \times 6.022 \times 10^{23} \times [0.4518 \times 10^{-7}]^2}{4}$$

$$= 1.057 \times 10^2 = 105.7$$
 gram/mole ≈ 106 g/mol

54. (7):
$$N_2O_{5(g)} \rightarrow 2NO_{2(g)} + \frac{1}{2}O_{2(g)}$$

Initial $a = 2.4 \times 10^{-2} M$ 0 0
After 1 hour $(a - x) = 1.6 \times 10^{-2} M$ x x
 $K = \frac{1}{t} \ln\left(\frac{a}{a - x}\right)$
 $k = \frac{2.303}{60} \log\left(\frac{2.4 \times 10^{-2}}{1.6 \times 10^{-2}}\right)$
 $k = \frac{2.303}{60} \log\left(\frac{3}{2}\right)$

 $k = 0.0067 = 6.7 \times 10^{-3} \text{ min}^{-1}$ or $k \approx 7 \times 10^{-3} \text{ min}^{-1}$. **55.** (78) : $C_6H_6 \xrightarrow{\text{Methylation}} C_6H_5CH_3$ 1 mol of benzene gives 1 mol of toluene. 78 g of benzene gives 92 g of toluene.

$$(W_{\text{theoretical}}) = \frac{10}{78} \times 92$$

% yield = $\frac{W_{\text{actual}}}{W_{\text{theoretical}}} \times 100 = \left[\frac{9.2}{10 \times 92} \times 78\right] \times 100 = 78\%$
56. (3) : For the given cell:
 $E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.059}{2} \log \left\{\frac{0.220}{(10^{-3})^2}\right\}$
= 2.97 - 0.0295 $\log[2.5 \times 10^5]$
= 2.97 - 0.0295 [5 + log 2.5] = 2.8 = 3 V
57. (21) : C(graphite) + $O_2 \longrightarrow CO_{2(g)}$
12 gram $\Delta H = -2.48 \times 10^2 \text{ kJ/mole}$
Total heat released by 1 gram = 2.48 $\times \frac{1}{12} \times 10^2$
= 20.67 kJ = 21 kJ
58. (12) : $_{23}\text{V} = 1s^22s^22p^63s^23p^63d^34s^2$
59. (6):
 $C - C - C - C = C$ (Pent-1-ene)
 $C - C - C = C$ (2-Methylbut-1-ene)
 $\frac{1}{C}$
 $C - C = C - C - C = (2-Methylbut-1-ene)$
 $\frac{1}{C}$
 $C - C = C - C - C - C (Pent-2-ene) \rightarrow 2 \text{ Geometrical Isomers}$
60. (2) : $K_p = K_c(RT)^{\Delta n_g}$
 $47.9 = K_c(0.083 \times 288)^1 \Rightarrow K_c = 2$
61. (c) : Given, $A = [a_{ij}]$ be a real matrix of order 3×3 such that $a_{i1} + a_{i2} + a_{i3} = 1$, for $i = 1, 2, 3$.
 $\therefore A = \begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{bmatrix} \Rightarrow A^3 = \begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{bmatrix}$
 \therefore Sum of all the entries of the matrix A^3 is $1 + 1 + 1 = 3$.
62. (c) : Four dice are thrown simultaneously.
 \therefore Total number of possible matrices $= 6 \times 6 \times 6 \times 6$
Now, let $A = \begin{bmatrix} a & b\\ c & d \end{bmatrix}$
If A is singular matrix $\Rightarrow |A| = 0$
 $\Rightarrow ad - bc = 0 \Rightarrow ad = bc$
 $\therefore a = 1, d = 6$ and $b = 3, c = 2 \rightarrow 4$ possible matrices
 $a = 2, d = 3$ and $b = 1, c = 6$

a = 3, d = 4 and b = 2, c = 6 \rightarrow 4 possible matrices Thus, total number of cases for singular matrix having different entries = 4 + 4 + 4 + 4 = 16: Total number of non-singular matrices having different entries = $6 \times 5 \times 4 \times 3 - 16 = 344$ The probability that the matrices are non-singular and have different entries = $\frac{344}{1296} = \frac{43}{162}$ 63. (b): We have, $\sin^7 x + \cos^7 x = 1$, $x \in [0, 4\pi]$ If $\sin x \neq 0$ and $\cos x \neq 0$, then $\sin^7 x < \sin^2 x$...(i) and $\cos^7 x < \cos^2 x$...(ii) Adding both (i) and (ii), we get $\sin^7 x + \cos^7 x < \sin^2 x + \cos^2 x$ $\Rightarrow \sin^7 x + \cos^7 x < 1$ Hence, $\sin^7 x + \cos^7 x = 1$ is the only possibility, when $\sin x = 1$ and $\cos x = 0$ or $\cos x = 1$ and $\sin x = 0$. $\therefore x = 0, \frac{\pi}{2}, \frac{5\pi}{2}, 2\pi, 4\pi \text{ [As } x \in [0, 4\pi]\text{]}$ \therefore Number of solutions = 5 **64.** (c) : (a) $[\vec{a}\vec{b}\vec{c}] + [\vec{c}\vec{a}\vec{b}] = 2[\vec{a}\vec{b}\vec{c}] = 2(\vec{a}\cdot(\vec{b}\times\vec{c}))$ $= 2\vec{a}\cdot\vec{a} = 2|\vec{a}|^2 = 8$ (b) Projection of \vec{a} on $\vec{b} \times \vec{c} = \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{|\vec{b} \times \vec{c}|} = \frac{\vec{a} \cdot \vec{a}}{|\vec{a}|} = |\vec{a}| = 2$ (c) If $\vec{a} \vec{b} \vec{c}$ are mutually perpendicular vectors, then $|\vec{a} \times \vec{b}| = |\vec{c}| \Rightarrow |\vec{a}| |\vec{b}| = |\vec{c}| \Rightarrow |\vec{b}| = \frac{|\vec{c}|}{2}$ Also, $|\vec{b} \times \vec{c}| = |\vec{a}|$ \Rightarrow $|\vec{b}||\vec{c}| = 2 \Rightarrow |\vec{c}| = 2$ and $|\vec{b}| = 1$ $|3\vec{a} + \vec{b} - 2\vec{c}|^2 = (3\vec{a} + \vec{b} - 2\vec{c}) \cdot (3\vec{a} + \vec{b} - 2\vec{c})$ $=9|\vec{a}|^{2} + |\vec{b}|^{2} + 4|\vec{c}|^{2} = (9 \times 4) + 1 + (4 \times 4)$ = 36 + 1 + 16 = 53(d) $\vec{a} \times ((\vec{b} + \vec{c}) \times (\vec{b} - \vec{c}))$ $=\vec{a}\times((-\vec{b}\times\vec{c})+(\vec{c}\times\vec{b}))=-2(\vec{a}\times(\vec{b}\times\vec{c}))=-2(\vec{a}\times\vec{a})=\vec{0}$ **65.** (d): Given, equation is $z^2 + 3\overline{z} = 0$ Let z = x + iy, then $\overline{z} = x - iy$ $\therefore \quad z^2 + 3\,\overline{z} = 0 \Longrightarrow (x + iy)^2 + 3(x - iy) = 0$ $\Rightarrow x^2 - y^2 + 2ixy + 3x - 3iy = 0$ $\Rightarrow x^2 - y^2 + 3x + i(2xy - 3y) = 0$ $\Rightarrow x^2 - y^2 + 3x = 0 \text{ and } y(2x - 3) = 0$...(i) When y = 0, $x^2 - y^2 + 3x = 0 \implies x = 0$ or x = -3When $y \neq 0 \Rightarrow x = \frac{3}{2}$; $x^2 - y^2 + 3x = 0 \Rightarrow y = \pm \frac{3\sqrt{3}}{2}$ \therefore Solutions are (0, 0), (-3, 0), $\left(\frac{3}{2}, \frac{3\sqrt{2}}{2}\right)$ and $\left(\frac{3}{2}, \frac{-3\sqrt{2}}{2}\right)$ \Rightarrow Total solutions = 4 \Rightarrow *n* = 4 $\therefore \quad \sum_{k=0}^{\infty} \frac{1}{4^k} = \frac{1}{1 - \frac{1}{4}} = \frac{4}{3}$ (Sum of infinite G.P.)

___unacademy

66. (d) : Let *L* be the line of intersection of planes $\pi_1: \vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 2$ and $\pi_2: \vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 2$ and let it cuts *xy* – plane at point *A*.

 \Rightarrow x - y = 2 and 2x + y = 2 [Using equations of given plane]

$$\Rightarrow x = \frac{4}{3}, y = \frac{-2}{3}$$
$$\therefore A\left(\frac{4}{3}, \frac{-2}{3}, 0\right)$$

Vector parallel to the line of intersection is given by

$$\vec{n} = \vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 2 & 1 & -1 \end{vmatrix} = -\hat{i} + \hat{j} + \hat{k}$$

Thus, equation of line of intersection is

$$\frac{x-\frac{4}{3}}{-1} = \frac{y+\frac{2}{3}}{5} = \frac{z-0}{3} = \lambda(\text{say})$$

Let coordinates of foot of perpendicular be

$$B\left(-\lambda + \frac{4}{3}, 5\lambda - \frac{2}{3}, 3\lambda\right)$$

$$\therefore \quad C(1, 2, 0) \text{ lies on the perpendicular line}$$

$$\therefore \quad \overline{CB} = \left(-\lambda + \frac{1}{3}\right)\hat{i} + \left(5\lambda - \frac{8}{3}\right)\hat{j} + (3\lambda)\hat{k}$$

Now, $\overrightarrow{CB} \cdot \overrightarrow{n} = 0$ $\Rightarrow \lambda - \frac{1}{2} + 25\lambda - \frac{40}{2} + 9\lambda = 0 \Rightarrow 35\lambda = \frac{41}{3} \Rightarrow \lambda = \frac{41}{105}$ Now, as $\alpha = -\lambda + \frac{4}{3}$, $\beta = 5\lambda - \frac{2}{3}$ and $\gamma = 3\lambda$ $\therefore \alpha + \beta + \gamma = 7\lambda + \frac{2}{3} = 7\left(\frac{41}{105}\right) + \frac{2}{3} = \frac{51}{15}$ \Rightarrow 35(α + β + γ) = $\frac{51}{15} \times 35 = 119$. **67.** (d): (a) $(p \Rightarrow \neg q) \lor (\neg q \Rightarrow p)$ $\equiv (\sim p \lor \sim q) \lor (q \lor p) \equiv (\sim p \lor p) \lor (\sim q \lor q) \equiv t \lor t \equiv t$ (b) $(p \Rightarrow q) \lor (\sim q \Rightarrow p)$ $\equiv (\neg p \lor q) \lor (q \lor p) \equiv (\neg p \lor p) \lor q \equiv t \lor q \equiv t$ (c) $(q \Rightarrow p) \lor (\sim q \Rightarrow p)$ $\equiv (\neg q \lor p) \lor (q \lor p) \equiv (\neg q \lor q) \lor p \equiv t \lor p \equiv t$ (d) $(\sim p \Rightarrow q) \lor (\sim q \Rightarrow p) \equiv (p \lor q) \lor (q \lor p)$ $\equiv (p \lor p) \lor (q \lor q) \equiv p \lor q$, which is not a tautology. **68.** (b): $[e^x]^2 + [e^x + 1] - 3 = 0$ $\Rightarrow [e^x]^2 + [e^x] - 2 = 0$ Let $[e^x] = y$, then we have $y^2 + y - 2 = 0$ $\Rightarrow (y+2)(y-1) = 0$ $\Rightarrow y = -2, 1$ \therefore $[e^x] = 1, -2 \Rightarrow [e^x] = 1 \Rightarrow x \in [0, \log_e 2)$ **69.** (d) : We have, $E_1: \frac{x^2}{x^2} + \frac{y^2}{x^2} = 1, a > b$ Also, it is given that, eccentricities $e_1 = e_2$.

 $\Rightarrow 1 - \frac{b^2}{a^2} = 1 - \frac{a^2}{B^2} (\because B > a)$ I(0,B)(0, b) $\Rightarrow B^2 = \frac{a^4}{a^2}$ (a,0) x (-a,0)(-ae,0)(ae,0) $\Rightarrow B = \frac{a^2}{b}$...(i) Now, as foci of E_2 are end (0, -B)points of minor axis of E_1 , therefore $B \cdot e = b$...(ii) From (i) and (ii), we get $e = \frac{b^2}{2}$ Since, $e^2 = 1 - \frac{b^2}{2}$ $\therefore e^2 = 1 - e$ $\Rightarrow e^2 + e - 1 = 0 \Rightarrow e = \frac{\sqrt{5} - 1}{2}.$ 70. (d): We have, $f(x) = \frac{\cos^{-1}\sqrt{x^2 - x + 1}}{\sqrt{\sin^{-1}\left(\frac{2x - 1}{2}\right)}}$ Clearly, f(x) will be well-defined if $1 \ge x^2 - x + 1 \ge 0$ and $0 < \frac{2x-1}{2} \le 1$ \Rightarrow $x(x-1) \leq 0$ and $1 < 2x \leq 3$ $\Rightarrow x \in [0,1] \cap x \in \left(\frac{1}{2}, \frac{3}{2}\right)$ $\Rightarrow x \in [0,1] \cap \left(\frac{1}{2}, \frac{3}{2}\right] \Rightarrow x \in \left(\frac{1}{2}, 1\right)$ $\Rightarrow \alpha = \frac{1}{2} \text{ and } \beta = 1 \quad \therefore \quad \alpha + \beta = \frac{3}{2}.$ 71. (d): We have, $f(x) = \begin{cases} \frac{x^3}{(1 - \cos 2x)^2} \log_e \left(\frac{1 + 2xe^{-2x}}{(1 - xe^{-x})^2} \right) , x \neq 0 \end{cases}$ x = 0Now, as *f* is continuous at x = 0 $\therefore \lim_{x \to 0} f(x) = \alpha$ Consider, $\lim_{x \to 0} f(x) = \lim_{x \to 0} \frac{x^3}{(1 - \cos 2x)^2} \log \left(\frac{1 + 2xe^{-2x}}{(1 - xe^{-x})^2} \right)$ $= \lim_{x \to 0} \frac{x^3 \times x}{(1 - 1 + 2\sin^2 x)^2} \frac{\log(1 + 2xe^{-2x}) - 2\log(1 - xe^{-x})}{x}$ $= \lim_{x \to 0} \frac{x^4}{4\sin^4 x} \cdot \frac{\log(1 + 2xe^{-2x}) - 2\log(1 - xe^{-x})}{x}$ $= \lim_{x \to 0} \left| \frac{1}{4} \left(\frac{2(xe^{-2x}(-2) + e^{-2x})}{(1+2xe^{-2x})} - \frac{2(xe^{-x} - e^{-x})}{(1-xe^{-x})} \right) \right| = \frac{1}{4} \times 4$ $\therefore \alpha = 1.$

$$\Rightarrow \frac{9}{4} < \frac{C}{36} \text{ and } \frac{100}{36} < \frac{C}{36}$$

$$\Rightarrow C > \frac{9 \times 36}{4} \text{ and } C > \frac{36 \times 100}{36}$$

$$\Rightarrow C > 81 \quad ...(i) \text{ and } C > 100 \qquad ...(ii)$$
Now, as $g^2 + f^2 - C > 0$

$$\therefore \left(-\frac{3}{2}\right)^2 + \left(\frac{5}{3}\right)^2 - \frac{C}{36} > 0$$

$$\Rightarrow \frac{C}{36} < \frac{9}{4} + \frac{25}{9} \Rightarrow C < 181 \qquad ...(iii)$$
Intersection point of $2x - y = 5$ and $x - 2y = 4$ is $(2, -1)$
and $(2, -1)$ lies inside the circle. So, $S(2, -1) < 0$

$$\therefore 36(2)^2 + 36(-1)^2 - 108(2) + 120(-1) + C < 0$$

$$\Rightarrow C < 156 \qquad ...(iv)$$

:. By (ii), (iii) and (iv), we have

$$100 < C < 156.$$

75. (a) : Let
$$L_1: \frac{(x-1)}{2} = \frac{(y-2)}{1} = \frac{(z-1)}{3}; \ \vec{r}_1 = 2\hat{i} + \hat{j} + 3\hat{k}$$

 $L_2: \frac{(x-2)}{1} = \frac{(y-\lambda)}{2} = \frac{(z-3)}{4}; \ \vec{r}_2 = \hat{i} + 2\hat{j} + 4\hat{k}$

est distance = Projection of \vec{a} on $\vec{r_1} \times \vec{r_2}$

$$= \frac{|a \cdot (\vec{r}_{1} \times \vec{r}_{2})|}{|\vec{r}_{1} \times \vec{r}_{2}|}$$

$$= \frac{|a \cdot (\vec{r}_{1} \times \vec{r}_{2})|}{|\vec{r}_{1} \times \vec{r}_{2}|}$$

$$= \frac{|a \cdot (\vec{r}_{1} \times \vec{r}_{2})|}{|\vec{r}_{1} \times \vec{r}_{2}|}$$

and
$$|\vec{r}_1 \times \vec{r}_2| = \sqrt{38}$$

 $\Rightarrow \frac{1}{\sqrt{38}} = \frac{|14 - 5\lambda|}{\sqrt{38}}$
 $\Rightarrow |14 - 5\lambda| = 1$
 $\Rightarrow 14 - 5\lambda = 1 \text{ or } 14 - 5\lambda = -1$
 $\therefore \lambda = \frac{13}{3} \text{ or } \lambda = 3$

$$\therefore \quad \lambda = \frac{1}{5} \text{ or } \lambda = 3$$

$$\therefore \quad \text{Integral value of } \lambda = 3.$$

76. (d): Given, vector
$$\vec{a}$$
 is coplanar with vector
 $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - \hat{j} + \hat{k}$. Also, we have \vec{a} is
perpendicular to $\vec{a} = 3\hat{i} + 2\hat{j} + 6\hat{k}$ and $|\vec{a}| = \sqrt{10}$.
 $\therefore \vec{a} = \lambda \hat{b} + \mu \hat{c} = \lambda(2\hat{i} + \hat{j} + \hat{k}) + \mu(\hat{i} - \hat{j} + \hat{k})$
 $= \hat{i}(2\lambda + \mu) + \hat{j}(\lambda - \mu) + \hat{k}(\lambda + \mu) = 0$
 $\Rightarrow 3(2\lambda + \mu) + 2(\lambda - \mu) + \hat{k}(\lambda + \mu) = 0$
 $\Rightarrow 3(2\lambda + \mu) + 2(\lambda - \mu) + \hat{k}(\lambda + \mu) = 0$
 $\Rightarrow 14\lambda + 7\mu = 0 \Rightarrow \mu = -2\lambda$
 $\Rightarrow \vec{a} = (0\hat{i}^2 + 3\hat{\lambda}\hat{j} + (-\lambda)\hat{k} \Rightarrow |\vec{a}| = \sqrt{10} |\lambda| = \sqrt{10}$
 $\Rightarrow |\lambda| + 1 = \lambda = 1 \text{ or } -1$
Now, as $|\vec{a} \cdot \vec{b}| = 1 \text{ or } -1$
Now, as $|\vec{a} \cdot \vec{b}| = 1 \text{ or } -1$
 $\Rightarrow -3\lambda(12) - \lambda(\hat{6}) = -42\lambda = 42 \text{ or } -42$.
Thus, the possible value is -42.
Thus, the possible value is -42.
Thus, the possible value is -42.
 $\Rightarrow -3\lambda(12) - \lambda(\hat{6}) = -42\lambda = 42 \text{ or } -42$.
Thus, the possible value is -42.
 $\Rightarrow \frac{dy}{dx} + 2\sin^2 x = 1 + y\cos 2x$ (a) $\frac{2}{3} \cdot 2 = \hat{6}$
 $\Rightarrow -3\lambda(12) - \lambda(\hat{6}) = -42\lambda = 42 \text{ or } -42$.
Thus, the possible value is -42.
 $\Rightarrow \frac{dy}{dx} + (-\cos 2x)y = \cos 2x$
 $\Rightarrow \frac{dy}{dx} + 2\sin^2 x = 1 + y\cos 2x$ (a) $\frac{\sin 2x}{2}$
Now, solution of differential equation is
 $y(\frac{\sin 2x}{2}) = -\frac{\sin 2x}{2} + e^{\frac{1}{2}}$.
Now, solution of differential equation is
 $y(\frac{\sin 2x}{2}) = -\frac{\sin 2x}{2} + e^{\frac{1}{2}}$.
Now, $y(0) = -1 + e^{-1/2} \Rightarrow (y(0) + 1)^2 = e^{-1}$
78. (a) : We have, $f(x) = \begin{cases} -\frac{4}{3}x^3 + 2x^2 + 3x, x > 0 \\ 3xe^k + e^k$], $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}(3x^2) + 4x + 3, x > 0 \\ 3xe^k + e^k$], $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + e^k$], $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + 3e^k$, $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + 3e^k$, $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + 3e^k$, $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + 3e^k$, $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x^2 + 4x + 3, x > 0 \\ 3xe^k + 3e^k$, $x < 0$
 $\Rightarrow f'(x) = \begin{cases} -\frac{4}{3}x$

.). interval $\left(-1, \frac{3}{2}\right)$. ations is $2y + \lambda z = \mu$ to above system of linear $B \neq O.$ (5) = 0= 140 ...(i) +4d = 56...(ii) d d = 102a + 5d] 75 d $\times 5 \times 17$ 3, 5 and 17 + 99) - (3 + 9 + 15 + 5 + 85 + 95) - (17) = 1251the set $2040 \text{ is } 1\} = 1251$

82. (4) : We have a function
$$f: R \to R$$
 defined as

$$f(x) = \begin{cases} 3\left(1 - \frac{|x|}{2}\right), & \text{if } |x| \le 2\\ 0, & \text{if } |x| > 2 \end{cases}$$
$$f(x+2) = \begin{cases} 3\left(\frac{4+x}{2}\right), & \text{if } -4 \le x \le -2\\ \frac{-3x}{2}, & \text{if } -2 < x \le 0\\ 0, & \text{if } x \in (-\infty, -4) \cup (0, \infty) \end{cases}$$

$$-f(x-2) = \begin{cases} \frac{-3x}{2} &, \text{ if } 0 \le x \le 2\\ \frac{3(x-4)}{2}, & \text{if } 2 < x \le 4\\ 0 &, & \text{if } x \in (-\infty, 0) \cup (4, \infty) \end{cases}$$
Now, $g(x) = f(x+2) - f(x-2)$

$$(1 + 2) - f(x-2)$$

$$(1 + 3) - f(x-2)$$

83. (2) : Region bounded by the given curves is the shaded region, shown below.



84. (8) : Given, constant term in the binomial expansion of $\left(2x^r + \frac{1}{x^2}\right)^{10} = 180$

Let the $(k + 1)^{\text{th}}$ term is $T_{k+1} = {}^{10}C_k (2x^r)^{10-k} \left(\frac{1}{x^2}\right)^k$ $= {}^{10}C_k(2)^{(10-k)} \cdot x^{r(10-k)} \cdot (x^{-2})^k = {}^{10}C_k(2)^{(10-k)}x^{10r-rk-2k}$ Now, for constant term 10r - rk - 2k = 0 $\Rightarrow r = \frac{2k}{10-k}$ and ${}^{10}C_k(2)^{10-k} = 180$ $\Rightarrow k=8$ $\therefore r=\frac{2\times 8}{10-8}=8.$ **85.** (720): Let $A = \{0, 1, 2, 3, 4, 5, 6, 7\}$. To find the number of bijective functions $f: A \to A$ such that f(1) + f(2) + f(3) = 3, the only possibility for this sum is 0 + 1 + 2 = 3. \Rightarrow Elements 1, 2, 3 in the domain can be mapped with 0, 1, 2 only. Therefore, number of bijective functions = $3! \times 5! = 6 \times 120$ = 720.86. (4): Given D.E. is $\left| (x+2)e^{\left(\frac{y+1}{x+2}\right)} + (y+1) \right| dx = (x+2)dy$, y(1) = 1.Let $y + 1 = Y \implies dy = dY$ and $x + 2 = X \implies dx = dX$ $\Rightarrow \left(Xe^{\frac{Y}{X}} + Y\right)dX = XdY \Rightarrow XdY - YdX = Xe^{\frac{Y}{X}}dX$ $\Rightarrow d\left(\frac{Y}{Y}\right)e^{\frac{-Y}{X}} = \frac{dX}{Y} \Rightarrow -e^{-\frac{Y}{X}} = \log |X| + C$ At y(1) = 1, X = 3, Y = 2 $\therefore \quad Y(3) = 2 \implies -e^{-\frac{2}{3}} = \log|3| + C$ $\Rightarrow e^{-\frac{Y}{X}} = e^{-\frac{2}{3}} + \log 3 - \log |X| > 0$ $\Rightarrow \log |X| < e^{-\frac{2}{3}} + \log_3 = \lambda \text{ (say)}$ $\Rightarrow |x+2| < e^{\lambda}$ $\Rightarrow -e^{\lambda} < x + 2 < e^{\lambda} \Rightarrow -e^{\lambda} - 2 < x < e^{\lambda} - 2$ $\Rightarrow \alpha + \beta = -4 \Rightarrow |\alpha + \beta| = 4$ 87. (96): We have digits 0, 2, 4, 6, 8. В CD Ε Α Number of possibility for A = 4Number of possibility for B = 4Number of possibility for C = 3Number of possibility for D = 2Number of possibility for E = 1

:. Number of numbers greater than 10,000 formed $= 4 \times 4 \times 3 \times 2 \times 1 = 96$.



88. (96) : Let
$$f(n) = \left(\frac{10}{11}\right)^n + \left(\frac{9}{11}\right)^n$$
, then
 $f'(n) = \left(\frac{10}{11}\right)^n \log\left(\frac{10}{11}\right) + \left(\frac{9}{11}\right)^n \log\left(\frac{9}{11}\right) < 0$
 $\Rightarrow f(n)$ is decreasing function.
Putting $n = 1, 2, 3, 4, \dots, we see that$
 $f(1) = \frac{10}{11} + \frac{9}{11} = \frac{19}{11} > 1$
 $f(2) = \left(\frac{10}{11}\right)^2 + \left(\frac{9}{11}\right)^2 = \frac{100}{121} + \frac{81}{121} = \frac{181}{121} > 1$
Similarly, $f(3)$ and $f(4) > 0$. But $f(5) = \left(\frac{10}{110}\right)^5 + \left(\frac{9}{11}\right)^5 < 1$
Similarly, for $n = 6, 7, 8, \dots, 100, f(n) < 1$.
 \therefore Required number of elements = 96.
89. (4):
 $\boxed{\frac{Class}{12.18} \frac{Frequency}{12.18} \frac{x_i}{12.1} \frac{15}{180} \frac{189}{12.18} \frac{12}{12} \frac{15}{180} \frac{189}{12.24} \frac{9}{21} \frac{115}{189} \frac{189}{22} \frac{133}{189} \frac{19}{2430} \frac{5}{5} \frac{277}{135} \frac{133}{189} \frac{19}{2430} \frac{5}{5} \frac{277}{135} \frac{135}{189} \frac{13}{1824} \frac{49}{189} \frac{9}{22}$ (Given)
 \therefore Mean $= \frac{3a + 9b + 504}{a + b + 26} = \frac{309}{22}$ (Given)
 $\Rightarrow 66a + 198b + 11088 = 309a + 309b + 8034 \dots (i)$
 $\Rightarrow 81a + 37b = 1018 \dots (i)$
 $Clearly, median = 12 + \frac{a + b + 26}{2} - (a + b)}{\frac{a + b + 26}{2} - (a + b)} \times 6 = 14$ (Given)
 $\Rightarrow \frac{a + b + 18}{2} - (a + b) = 2$
 $\Rightarrow a + b = 18 \dots (i)$
 $By (i)$ and (ii), we get $a = 8, b = 10$
 $\therefore (a - b)^2 = (8 - 10)^2 = 4$
90. (3125) : Here, $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
 $b = 3 \times 3$ matrix such that $AB = BA$
 $\Rightarrow \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} p & q & r \\ s & t & u \\ v & w & x \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
 $\Rightarrow \frac{s}{s} = q, t = p, u = r, v = w$
So, we have to select 5 elements for B.
 \therefore Total number of ways $= 5^5 = 3125$.

