# **Mole Concept**

# CHEMISTRY

• Chemistry is defined as the study of the composition, properties and interaction of matter.

# **Branches of Chemistry**

# (1) Physical Chemistry

The discipline of chemistry concerned with the way in which physical properties of substances depend on and influence their chemical structure, properties, and reactions.

# (2) Inorganic Chemistry

The discipline of chemistry in which structure, composition, and behavior of inorganic compounds. All the substances other than the carbon-hydrogen compounds are classified under the group of inorganic substances.

# (3) Organic Chemistry

The branch in which the study of the structure, composition and the chemical properties of organic compounds is known as organic chemistry.

#### (4) Biochemistry

The discipline which deals with the structure and behavior of the components of cells and the chemical processes in living beings is known as biochemistry

# (5) Analytical Chemistry

The branch of chemistry dealing with separation, identification and quantitative determination of the compositions of different substances

#### MATTER

 It exists in three physical states, e.g., Solid, Liquid and Gas. It also has two other states namely Bose-Eistein condensite and Plasma

# Concept Ladder



Chemistry, from the ancient Egyptian word "khēmia" meaning transmutation of earth, is the science of matter at the atomic to molecular scale, dealing primarily with collections of atoms, such as molecules, crystals, and metals.

# **Rack your Brain**



How was mass of single carbon aotm determined?

#### **Concept Ladder**



Antoine Lavoisier is known as father of chemistry. He developed an experimentally based theory of the chemical reactivity of oxygen and coauthored the modern system for naming chemical substances.

# Definition

**Matter** is anything that occupies mass and space.



# **Characteristics**

- (1) Solids have definite volume and definite shape.
- (2) Liquids have definite volume but definite shape. They take the shape of the container in which they are kept.
- (3) Gases alve neither definite volume nor definite shape. They occupy completelty the container in which they are kept.

These three states are interconvertible on changing the conditions of temperature and pressure.

# **Classification of Matter On the Basis of Purity**

Matter can be classified broadly as mixture or pure substances, which can be further subdivided as shows below.

#### **Concept Ladder**



and characteristic properties. It cannot be separated into components by physical separation methods, i.e. without breaking chemical bonds. They can be solids, liquids or gases.



Mole Concept

#### (1) Mixture

Generally pure substances are added together to form a mixture. Also, a mixture can be obtained by mixing two mixtures. For example, sugar solution in water, air, tea etc.

# (2) Homogenous Mixture

A mixture in which the components completely mix with each other and its composition is uniform throughout. For example, salt solution, sugar solution, air etc.

# (3) Heterogenous Mixture

A mixture which doesn't have same composition throughout and different components sometimes can be observed. For example, grains and pulses along with some dirt (often stone) pieces , mixture of salt and sugar etc.

#### (4) Pure Substances

They have fixed composition, whereas mixture may contain the components in any ratio and its composition is variable. For example, gold, silver, copper, water, glucose etc.

#### (5) Element

Element has only one type of particles, atoms or molecules. For example silver, copper, sodium, hydrogen, oxygen etc.

### (6) Molecule

When two or more atoms combine molecule is formed. Hydrogen, oxygen, and nitrogen gases consists of molecules in which atoms of same elements combine to give their respective molecules.

### (7) Compound

Compounds are always formed when substances combine in different ratios by mass. For example water ammonia, carbon

### Definition

A **mixture** contains two or more substances present in it in any ratio which are called its components.

# **Concept Ladder**

Alloys are mixtures of two or more metals or a metal and a non-metal and cannot be separated into their components by physical methods. For example, brass is a mixture of approximately 30% zinc and 70% copper.

# **Rack your Brain**



How do we judge whether milk, ghee, butter, salt, spices, mineral water or juice that we buy from the market are pure?

# Definition

When two or more atoms of different elements combine, the molecule of a compound is formed.

# **STATES OF MATTER**



monoxide, sodium chloride etc.

### **Properties of Matter and Their Measurements**

 The properties of a substance have unique characteristics and are classified into physical and chemical properties.

# (1) Physical Properties

Those properties which can be measured or observed without changing the identity or composition of the substance are known as physical properties, .e.g, colour, melting point, boiling point, odour etc.

# (2) Chemical Properties

Those properties which describe a matter's 'potential' to undergo some chemical changes are konwn as chemical properties, e.g., characteristics reactions of different substances which include acidity or basicity, combustibility etc.

### (3) Measurement

Any quantitative observation represented by a number followed by a unit in which it is measued is called measurement, such as length, are, volume etc.

# PHYSICAL QUANTITIES AND SI UNITS OF MEASUREMENT

 11<sup>th</sup> general conference of weights and measures recommended the use of international system of units in 1960. SI Units is abbreviated as (after the French expression La System International de units).

### **Fundamental Units**

• The SI system has seven basic/fundamental units of physical quantities as follows:

# Concept Ladder



Physical properties like melting and boiling points can be the result of the components present inside a system.

# **Rack your Brain**



How do we judge whether milk, ghee, butter, salt, spices, mineral water or juice that we buy from the market are pure?

# **Concept Ladder**



There are other unit systems used for measurement of different physical and chemical quantities For example, FPS (Foot, poundal, second), CGS (centimeter, gram, second).

# **Rack your Brain**



Can you guess what is BTU?

Physical Quantity	Abbreviation	Name of unit	Symbol
time	t	second	S
mass	m	kilogram	kg
length	l	metre	m
temperature	Т	kelvin	К
electric current	I	ampere	А
light intensity	lv	candela	Cd
amount of substance	n	mole	mol

# **Derived Units**

 The units obtained by combination of basic units are known as derived units e.g. velocity is expressed as distance/time. Hence unit is m/s or ms-1. Some common derived units are:

Concept	Ladder
---------	--------



Fundamental units are the basis of derived units.. Derived units are obtained when fundamental units are divided or multiplied together but cannot be obtained on addition or substraction.

Physical Quantity	Definition	SI Unit
volume	length cube	m <sup>3</sup>
area	length square	m²
speed	distance travelled	ms⁻¹ per unit time
acceleration	speed changed	ms <sup>-2</sup> per unit time

Fraction	Prefix	Symbol	Multiple	Prefix	Symbol
10 <sup>-1</sup>	deci	d	10 <sup>1</sup>	Deka	da
10 <sup>-2</sup>	centi	С	10 <sup>2</sup>	Hecta	h
10-3	milli	m	10 <sup>3</sup>	kilo	k
10 <sup>-6</sup>	micro	m	10 <sup>6</sup>	Mega	Μ
10 <sup>-9</sup>	nano	n	10 <sup>9</sup>	Giga	G
10 <sup>-12</sup>	pico	р	10 <sup>12</sup>	Tera	Т
10 <sup>-15</sup>	femto	f	10 <sup>15</sup>	Peta	Р
10 <sup>-18</sup>	atto	а	10 <sup>18</sup>	Exa	E

# **Standard Prefixes**

# **Precision and Accuracy**

# Precision

Precision is the closeness of various measurements done for the same quantity

# Accuracy

Accuracy is the agreement of a subjected value to the true value.

**Ex:** Let the true weight of a substance be 3.00g.

The measurement reported by three students are as follows:

# **Concept Ladder**



The accuracy of any measurment depends upon two parameters. First being the device and second being the skill of operator.

Sudent	Measure	Average/g	
	1	2	
А	2.95	2.93	2.94
В	3.01	2.99	3
С	2.94	3.05	2.99



POOR ACCURACY POOR PRECISION

# **Case of A student**

It is the case of precision but no accuracy since measurements one close but not accurate.

# Case of B student

Measurements are very close (precision) and accurate (Accuracy).

# **Case of C student**

Measurement is not close (no precision) and not accurate (no accuracy).

# SOME IMPORTANT DEFINITION

# **Mass and Weight**

- It is the quantity of matter present in it while weight is the force exerted by gravity on an object.
- The mass of a substance is constant. Weight varies from one place to another due to change in gravity.
- SI unit of mass is kg.

#### Volume

- Volume is often quantified numerically using the SI derived unit, the cubic meter (m<sup>3</sup>).
- Volume of liquids or solutions is measured by using burette, pipette, graduated cylinder, or volumetric flask.

# Density

- The mass occupied per unit volume of a substance is known as Density. The symbol most often used for density is ρ
- SI unit of density is kg m<sup>-3</sup>.
- Other units of density include lb ft<sup>-3</sup> and g cm<sup>-3</sup>.
- Density is usually calculated with respect to a standard substance. This is known as relative density.



GOOD ACCURACY GOOD PRECISION



# Definition

Volume is the quantity of threedimensional space enclosed by some closed boundary, for example, the space that a substance (solid, liquid, gas, or plasma) or shape occupies or contains.

# Temperature

- Temperature is a physical property of matter which tell us about the degree of heat content.
- There are three common scales for measurement of temperature — K (kelvin, °F (degree Fahrenheit) and)°C (degree Celsius).
- The temperature on two scales is related to each other by the following relationship: °F = 9/5(°C) + 32 K = °C + 273-15

# LAWS OF CHEMICAL COMBINATION

• The combination of elements to form compounds chemically is the result of the five basic laws:





# Law of Conservation of Mass

- The law states that matter can neither be created nor destroyed.
- This law was given by Antoine Lavoisier in 1789.



- **1** Five grams of  $KClO_3$  yid 3.04 1 g and 1.36 L of oxygen at standard temperature and pressure. Show that these figures support the law of conservation of mass within limits of ±0.4% error.
- A1 According to gram-molecular volume law, 22.4 L of all gases and vapours at STP weight equal to their molecular weights denoted in grams.

$$\therefore \text{ Weight of 1.36 L of oxygen at STP} = \frac{32 \times 1.36}{22.4} = 1.943 \text{ g}$$
  
Weight of KCl formed = 3.041 g (given)  
$$\therefore \text{ Total weight of product (KCl + O_2)} = 3.041 + 1.943 = 4.984 \text{ g}$$
  
Error = 5 - 4.984 = 0.001 g  
$$\therefore \text{ \% error} = \frac{0.016 \times 100}{5} = 0.32$$

Hence, the law of conservation of mass is valid within limits of -0.4% error. Thus, the law is supported.

# • He performed experimental studies for combustion reactions for reaching to the above conclusion.

• This law did form the basis for several later developments in chemistry. In fact, this was the result of absolutely exact measurement of masses of reactants and products, and carefully planned experiments done by Lavoisier.

# **Rack your Brain**

What will happen when hydrogen and sulphur combine in the ratio 1:16 by mass?

# Law of Definite Proportions

- This law was given by French chemist, Joseph Proust.
- It was stated by him that a given compound always contains the same proportion of elements by weight.
- Proust considered two samples of cupric carbonate — one of the samples from natural origin and the other was of synthetic.
- He found that the composition of elements present in it was similar for both the samples as shown below:

# Concept Ladder



Law of constant composition is not true for all types of compounds but true only for the compounds obtained from one isotope.

	% of copper	% of oxygen	% of carbon
Natural sample	51.35	9.74	38.91
Synthetic sample	51.35	9.74	38.91

 Thus, irrespective of the source, a given compound always contains same elements in the same proportion. The validity of this law has been confirmed by various experiments. It is sometimes also referred to as Law of definite composition.

# **Rack your Brain**

Can we apply the law of definite proportion in case of nonstoichiometric compounds and polymers?

- Q2 0.7 g of iron reacts directly with 0.4 g of sulphur to form ferrous sulphide. If 2.8 g of iron is dissolved in dilute HCl and excess of sodium sulphide solution is added, 4.4 g of iron sulphide is precipitated. Prove the law of constant composition.
- A2 The ration of the weight of iron and sulphur in the first sample of the compound is Fe : S :: 0.7 : 0.4 or 7 : 4. According to the second experiment, 2.8 g of iron gives 4.4 g ferrous sulphide, or 2.8 g Fe combines with S = 4.4 - 2.8 = 1.6 g Therefore, the ratio of the weights of Fe : S :: 2.8 : 1.6 or 7.8. Since the ratio of the weights of the two elements is same in both the cases, the law of constant composition is true.

### **Law of Multiple Proportions**

- The law was proposed by Dalton in 1803.
- According to this law, when two elements combine to form more than one compound, the mass of one element that combines with fixed mass of the other element, are in the ratio of simplest whole numbers.
- For example,  $H_2$  combines with  $O_2$  to form two compounds, namely, water and  $H_2O_2$ .

Hydrogen	+	Oxygen	$\longrightarrow$	Water
2g		16g		18g
Hydrogen Peroxide	+	Oxygen	$\longrightarrow$	Hydrogen
2g		32g		34g

# **Concept Ladder**

The law, which was based on Dalton's observations of the reactions of atmospheric gases, states that when elements form compounds, the proportions of the elements in those chemical compounds can be expressed in small whole number ratios.

- Elements X and Y form two different compounds. In the first compound, 0.324 g X is combined with 0.471 g Y. In the second compounds, 0.117 g X is combined with 0.509 g Y. Show that these data illustrate the law of multiple proportions.
- A3 In the first compound 0.324 g of X combines with 0.471 g of Y. In the second compound 0.117 g of X combines with 0.509 g of Y.

Therefore, 0.324 g of X combines with the weight of Y =  $\frac{0.509 \times 0.324}{0.117}$  = 1.4095 g

Now, the weights of Y that combine with the same weight of X, i.e., 0.324 g of it, are in the ratio of 0.471 : 1.4095 or 1 : 3. The ratio, being simple, illustrates the law of multiple proportions.

#### **Gay Lussac's Law of Gaseous Volumes**

- This law was given in 1808 by Gay Lussac.
- It was observed that when gases combine together in a chemical reaction they combine in a simple ratio by volume provided all the gases are at same pressure and temperature.
- 1000 mL of H<sub>2</sub> combine with 500 mL of O<sub>2</sub> to give 1000 mL of H<sub>2</sub>O vapour.

$H_2$	+	0 <sub>2</sub>	$\longrightarrow$	$H_2O$
1000 mL		500 mL		1000 mL

# **Previous Year's Question**

# 8

Equal masses of  $H_2$ ,  $O_2$  and methane have been taken in a container of volumes V at temperature 27°C in identical conditions. The ratio of the volumes of gases  $H_2:O_2$ :methane would be

[AIPMT-2014]

(1) 8 : 16 : 1	(2) 16 : 8 : 1
(3) 16 : 1 : 2	(4) 8 : 1 : 2



- Q4 Air contains 21% oxygen by volume. Calculate the theroretical volume of air which will be required for burning completely 500 cubic ft of acetylene gas  $(C_2H_2)$ .
- $\begin{array}{ccc} \textbf{A4} & 2\text{C}_{2}\text{H}_{2} + 5\text{O}_{2} \\ & & 2 \text{ vol} \end{array} \xrightarrow{5 \text{ vol}} 5 \text{ vol} \xrightarrow{4 \text{CO}_{2}} + 2\text{H}_{2}\text{O}\left(\text{steam}\right) \end{array}$

According to the above equation: 2 volumes of acetylene require 5 volumes of  $O_2$  for combustion. 5 × 500 1050 cm ft

 $\therefore$  500 eu. ft. of acetylene will require  $O_2 = \frac{5 \times 500}{2} = 1250$  cu. ft.

Hence, the quantity of air required  $=\frac{100 \times 1250}{21} = 5952$  cu.ft.

# Avogadro law

 Avogadro proposed that when equal volumes of gases at same temperature and pressure they should contain equal number of molecules.

- Avogadro gave the distinction between atoms and molecules which is quite understandable in the present times.
- If we consider, reaction of H<sub>2</sub> and O<sub>2</sub> to produce H<sub>2</sub>O, we see that two volumes of H<sub>2</sub> combine with one volume of O<sub>2</sub> to give two volumes of H<sub>2</sub>O without leaving any unreacted O<sub>2</sub>.

# **DALTON'S THEORY OF ATOM**

# **Rack your Brain**

A container of 5 liter consists  $CO_2$  gas. Another container of half the volume consists  $O_2$  at same condition. Would it follow Avogadro's law?



J. Dalton gave his famous theory of atom in 1803. The main postulates of this theory

- Atom is considered as a hard, dense and\_ the smallest indivisible particle of matter.
- Each element consists of a identical kind of atoms.
- The properties of elements differ because of difference in the kinds of atoms contained in them.
- This theory provides a satisfactory basis for the law of chemical combination.

# **Limitations of Dalton's Theory**

• This theory fails to explain why the atoms of different kinds should differ in mass, valency etc.

# **Concept Ladder**

Dalton's atomic theory had certain explanation of Laws of Chemical Combination :

Law of conservation of mass : Matter consists of atoms and can neither be created nor destroyed.

Law of constant composition: When atoms of same or different elements combine together to form compounds, they combine in a fixed ratio, a simple whole number ratios.

- The discovery of isobars and isotopes demonstrated that atoms of same elements may have different atomic masses (isotopes) and atoms of different kinds may have same atomic masses (isobars).
- The discovery of various sub-aomic particles like electrons, protons, X-rays, etc. during the late 19th century lead to idea that atom was no longer an indivisible and the smallest particle of the matter.

# ΑΤΟΜ

- element is formed of smallest Each particles called 'ATOM'.
- Atom is derived from Greek language Atoms means 'Not to be cut'.

# **Atomic Mass**

Atomiic mass of an element can be defined by a number which indicates how many times the mass of one atom of the element is heavier in comparison to  $\frac{1}{12}$  th part of the

mass of one atom of Carbon-12.

Atomic mass = 
$$\frac{[Mass of an atom of the element]}{\frac{1}{12} \times [Mass of an atom of carbon-12]} = \frac{Mass of an atom in amu}{1 amu}$$

Definition

Atomic mass unit (amu) or Unified mass (u) • The quantity  $\frac{1}{12}$  × mass of an atom of C-12 is

∴ 1 a.m.u. = 
$$\frac{1.9924 \times 10^{-20}}{12}$$
 kg

The atomic mass as the weighted average mass of all naturally occurring isotopes of the element.

• A.A.M. = 
$$\left(\frac{\% \text{ abundance isotope 1}}{100}\right) \times (A_1) + \left(\frac{\% \text{ abundance isotope 2}}{100}\right) \times (A_2) + \dots$$

# **Rack your Brain**

Can you find law of multiple proportion and law of reciprocal proportion form Dalton's Atomic Theory Postulate?



Average atomic mass is defined as the average mass of isotopes of an element naturally present.



Calculate molecular mass of glucose (C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>) molecule.

Molecular mass of glucose (C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>) = 12(12.011 u) + 22(1.008 u) + 11(16.00 u) = 342.308 u

# Average Molecular Mass (AMM)

Average Molecular Mass

total mass total mole of molecules

 Let a sample contains n<sub>1</sub> mole of molecules with molecular mass M<sub>1</sub> and n<sub>2</sub> mole of molecules with molecualr mass M<sub>2</sub>, then

$$M_{av} = \frac{n_1 M_1 + n_2 M_2}{n_1 + n_2}$$

# **Formula Unit Mass**

- The mass of a formula unit present in ionic compound is known as formula unit mass.
- For example, the formula unit mass of sodium chloride (NaCl) is 58.5 u.

# **Gram Atomic Mass (GAM)**

- Gram atomic mass of an atom expressed in grams.
- Number of gram atoms

=  $rac{ ext{Mass of element in grams}}{ ext{GAM of element}}$ 

# Gram Molecular Mass (GMM)

- That amount of substance having mass in grams which is equal to its molecular mass of a substance expressed in grams is known as gram molecular mass.
- It is also called one gram molecule.
- Number of gram molecules
  - = Weight of substance GMM of substance

# **Previous Year's Question**



An element, X has the followign isotopic composition : <sup>200</sup>X:90% <sup>199</sup>X:8.0% <sup>202</sup>X:2.0%

# [AIPMT-2007]

(1) 201 amu (3) 199 amu (2) 202 amu(4) 200 amu

**Rack your Brain** 



There is a compound XCl<sub>2</sub> which is ionic in nature. Can this compound have molecular mass?

# Definition

The atomic mass of an element expressed in grams is called gram atomic mass.

Definition



The molecualr mass of a compound expressed in grams is called gram molecular mass.

#### **MOLE CONCEPT AND MOLAR MASSES**

- A mole is amount of a substance which consists of as many entities as there are atoms in 12 g (or 0.012 kg) of the C-12 isotope.
- The mass of a C-12 was calculated by a mass spectrometer and was found to be equal to 1.992648 × 10<sup>-23</sup> g.
- Number of atoms in one mole of carbon

= 
$$\frac{12 \text{ g mol}^{-1} \text{ of }^{12}\text{C}}{1.992648 \times 10^{-23} \text{ g} \left( {}^{12}\text{C atom} 
ight)^{-1}}$$

= 6.0221367 × 10<sup>23</sup> atoms mol<sup>-1</sup> ≈ 6.022 × 10<sup>23</sup> atoms mol<sup>-1</sup>

# **Molar Mass**

- The numercal value of molar mass of a substance in grams is numerically equal to atomic or molecular formula mass in u. Units of molar mass are g mol<sup>-1</sup> or kg mol<sup>-1</sup>.
- ∴ Molar mass of CO<sub>2</sub> = 12.011 + 2(16.0) = 44.011 g mol<sup>-1</sup>

Molar mass of NaCl = 23.0 + 35.5 = 58.5 g mol<sup>-1</sup>

# **Molar Volume**

- According to Avogadro's hypothesis, equal volumes of different gases under similar conditions of temperature and pressure contian equal number of molecules.
- A mole of a gas at STP (standard temperature and pressure), viz., 1 atm and 273 K (0°C) contains NA molecules (6.022 × 10<sup>23</sup>).
- 1 mole of a gas at 1 bar and 273 K consiting of N<sub>A</sub> molecules will have a volume of 22.7 liters which is considered to be **new STP** condition.

# **Previous Year's Question**

# 6

If Avogadro number  $N_A$ , is changed from 6.022 × 10<sup>23</sup> mol<sup>-1</sup> to 6.022 × 10<sup>20</sup> mol<sup>-1</sup>, this would change

#### [AIPMT-2015]

 (1) the mass of one mole of carbon
 (2) the ratio of chemical species to each other in a balanced equation
 (3) the ratio of chemical species to each other in a compound
 (4) the definition of mass in units of grams.

# **Definition**

substance in called its molar mass.





present in 1 cm<sup>3</sup> of an ideal gas at STO is called Loschdmidt number

 $=\frac{6.022\times10^{23}}{22400}$  mL = 2.69 × 10<sup>19</sup>

# **BASIC CONCEPTS OF ATOM**

# To calculate no. of p, n and e⁻

# (a) In case of neutral atom

Atom	р	e-	n
Carbon (C)	6	6	6
Nitrogen (N)	7	7	7

# (b) In case of ions

lon	р	e⁻	n
Oxide ion (O <sup>2–</sup> )	8	8 + 2 = 10	8
Nitride ion (N <sup>3-</sup> )	7	7 + 3 = 10	7

# (c) In case of molecule

# **Ex:** $CH_4$

Total number of atoms in  $CH_4 = 5$ 

Molecule	Element	р	e⁻	n
CH4	Carbon	6	6	6
	Hydrogen (1 × 4)	4	4	0
Total		10	10	6

**Ex:**  $N_2$ Total number of atoms in  $N_2 = 2$ 

Molecule	Element	р	e-	n	
N <sub>2</sub>	Nitrogen (1 × 2)	2 × 7 = 14	2 × 7 = 14	2 × 7 = 14	
Total		14	14	14	

# (d) In case of Charge on Molecule

**Ex:** (NH<sub>4</sub>)<sup>+</sup>

Total number of atoms in  $(NH_4)^+ = 5$ (No. of N-atoms = 1, No. of H-atoms = 4)

Molecule	Element	р	e-	n
(NH <sub>4</sub> ) <sup>+</sup>	Nitrogen	7	6	7
	Hydrogen (1 × 4)	4	4	0
Total		11	10	7

# **CALCULATION OF MOLES**



 Amount of substance which consist of Avogadro's number (6.022 × 10<sup>23</sup>) of atoms if the substance is atomic or Avogadro's number (6.022 × 10<sup>23</sup>) of molecules.

OR

In case of gaseous substance mole is the amount of gas which has a volume of 22.4 litres at STP.

# **Previous Year's Question**

Which has maximum molecules?

	[AIPMT]
(1) 7 g N <sub>2</sub>	(2) 2 g H <sub>2</sub>
(3) 16 g NO <sub>2</sub>	(4) 16 g O <sub>2</sub>



# Examples :

# (a) In case of atoms :

 1 Mole of carbon atoms = N<sub>A</sub> carbon atoms OR

1 Mole of carbon atoms =  $6.022 \times 10^{23}$  carbon atoms

• 1 Mole of nitrogen atoms =  $N_A$  nitrogen atoms =  $6.022 \times 10^{23}$  Nitrogen atoms

# (b) In case of ions :

- 1 Mole of O<sup>2-</sup> = N<sub>4</sub> O<sup>2-</sup> ions = 6.022 × 10<sup>23</sup> O<sup>2-</sup> ions
- 1 Mole of N<sup>3-</sup> = N<sub>A</sub> N<sup>3-</sup> ions = 6.022 × 10<sup>23</sup> N<sup>3-</sup> ions
- 1 Mole of  $NH_4^+ = N_A NH_4^+$  ions = 6.022 × 10<sup>23</sup>  $NH_4^+$  ions

# **Previous Year's Question**

Suppose the elements X and Y combine to form two compounds  $XY_2$  and  $X_3Y_2$ . When 0.1 mole of  $XY_2$  weighs 10 g and 0.05 mole of  $X_3Y_2$  weighs 9g, the atomci weights of X and Y are

# [NEET-2016]

(1) 40, 30	(2) 60, 40
(3) 20, 30	(4) 30, 20

# (c) In case of molecules :

- 1 Mole of N<sub>2</sub> = N<sub>A</sub> N<sub>2</sub> molecules = 6.022 × 10<sup>23</sup> N<sub>2</sub> molecules
- 1 Mole of O<sub>2</sub> = N<sub>A</sub> O<sub>2</sub><sup>-</sup> molecules = 6.022 × 10<sup>23</sup>
   O<sub>2</sub> molecules
- 1 Mole of  $CO_2 = N_A CO_2$  molecules = 6.022 ×  $10^{23} CO_2$  molecules

# (d) In case of Acid/Base/Salt/Double salts :

- 1 Mole of HCl = N<sub>A</sub> HCl = 6.022 × 10<sup>23</sup> HCl
- 1 Mole of NaOH =  $N_A$  NaOH = 6.022 × 10<sup>23</sup> NaOH
- 1 Mole of  $NH_4OH = N_A NH_4OH = 6.022 \times 10^{23}$  $NH_4OH$
- 1 Mole of NaCl =  $N_A$  NaCl = 6.022 × 10<sup>23</sup> NaCl
- 1 Mole of FeSO<sub>4</sub>.(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>.6H<sub>2</sub>O = 6.022 × 10<sup>23</sup> FeSO<sub>4</sub>.(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>.6H<sub>2</sub>O

# **Previous Year's Question**

The number of moles of oxygen in one litre of air containing 21% oxygen by volume, under standard conditions, is

[AIPMT]

- (1) 0.0093 mol
- (2) 2.10 mol
- (3) 0.186 mol
- (4) 0.21 mol

How many g atoms are there in on atom?

$$= 1.66 \times 10^{-24}$$
 g atom or mol

- From 200 mg of CO<sub>2</sub>, 10<sup>21</sup> molecules are removed. How many grams and moles of CO<sub>2</sub> are left.
- A9 44 g of CO<sub>2</sub> = 1 mol = 6.023 × 10<sup>23</sup> molecules  $\therefore 6.023 \times 10^{23}$  molecules = 44 g of CO<sub>2</sub> 10<sup>21</sup> molecules =  $\frac{40 \times 10^{21} \times 10^{3}}{6.023 \times 10^{23}}$ Weight of CO<sub>2</sub> left = 200 - 73.05 = 126.9 mg =  $\frac{126.9}{10^{3}}$  = 0.1269 g 44 g of CO<sub>2</sub> = 1 mol 0.1269 g of CO<sub>2</sub> =  $\frac{1}{44} \times 0.1269$  = 0.0028 mol



The volume of a drop of water is 0.04 mL. How many  $H_2O$  molecules are there in a drop of water? d = 1.0 g mL<sup>-1</sup>.

A13 Volume of 1 drop of  $H_2O = 0.04$  mL Weight of  $H_2O =$  Volume × Density = 0.04 × 1 = 0.04 g 1 mole of  $H_2O =$  18 g = 6.023 × 10<sup>23</sup> molecules

 $\therefore 0.04 \ g = \frac{6.023 \times 10^{23} \times 0.04}{18} = 1.3384 \times 10^{21} \text{ molecules}$ 

# **PRECENTAGE COMPOSITION**

• The percentage fo any element or constituent in a compound is the number of parts by mass of that element or constituent present in 100 parts by mass of the c

It is calculated as follows:

- First calculated the molecular mass of the compound from its formula by adding the atomic masses of the elements present in 100 parts by mass of the compound.
- Then calculate the percentage of the element or constituents by using the relation:

# **Previous Year's Question**

An organic Compound contains carbon, hydrogne and oxygen. its elemental analysis gave C, 38.71% and H, 9.67%. The empirical formula of the compound would be

[AIPMT-200	8]	
------------	----	--

(1) CHO	(2) CH <sub>4</sub> O
(3) CH <sub>3</sub> O	(4) CH <sub>2</sub> O

# Mass% of an element = $\frac{Mass of element \times 100 in the compound}{Molar mass of the compound}$

214 Calculate the percentage composition of various elements in the following compoud : Blue vitriol (CuSO<sub>4</sub>.5H<sub>2</sub>O).

A14 Molar mass of  $CuSO_4.5H_2O = 63.5 + 32 + 4 \times 16 + 5 \times 18 = 249.5$  g

Mass% of Cu 
$$= \frac{63.5 \times 100}{249.5} = 25.45\%$$

Mass% of S = 
$$\frac{32 \times 100}{249.5}$$
 = 12.82 %

Mass% of O = 
$$\frac{16 \times 9 \times 100}{249.5}$$
 = 57.71 %

Mass% of H = 
$$\frac{10 \times 1.008 \times 100}{249.5}$$
 = 4.040 %

# EMPIRICAL AND MOLECULAR FORMULAE Empirical Formula

- Empirical formula gives the simplest whole number ratio of the various atoms present in a compound.
- For example, the empirical formula of glucose is CH<sub>2</sub>O, that of benzene is CH, and that of hydrogen peroxide is OH.

# **Molecular Formula**

- It represents the exact number of different types of atoms present in a molecule of a compound.
- For example, molecular formual of glucose is C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>, that of hydrogen peroxide is H<sub>2</sub>O<sub>2</sub>.

# Relation Between Empirical and Molecular Formulae

• The molecualr formula of a compound is a simple whole number multiple of its empirical formula,

# Molecualr formual = n × Empirical formula

where n is any integer, e.g., 1, 2, 3, ..., etc. The value of n is obtained as

# n = <u>Molar mass</u> Empirical formula mass

# **Molar Mass of a Volatile Compound**

 It is determined by Victor Meyer's method. It is based on the principle that 22.4 L of vapours of a volatile compound at STP have mass equal to the gram molecualr mass or by the relation given below.

# Molar mass = 2 × Vapour density Calculation of the Empirical and Molecular Formulae

- Conversion of mass percent to grams.
- Convert into number of moles of each element.
- To calculate the simplest whole number.
- Writing empirical formula.
- Writing molecular formula.

# **Rack your Brain**

Can we apply percentage composition in case of nonstochiometric comopounds like Fe<sub>0.95</sub>O?



EMPIRICAL FORMULA



# Q15 A substance, on analysis, gave the following percentage composition: Na = 43.4%, C = 11.3% and O = 45.3%. Calculate the empirical formula. (Na = 23, C = 12, O = 16)

A15	Element	Percentage composition	Atomic ratio	Least ratio
	Sodium	43.4	$\frac{43.4}{23} = 1.89$	$\frac{1.89}{0.94} = 2$
	Carbon	11.3	$\frac{11.3}{12} = 0.94$	$\frac{0.94}{0.94} = 1$
	Oxygen	45.3	$\frac{45.3}{16} = 2.83$	$\frac{2.84}{0.94} = 3$

Hence, the empirical formula is  $Na_2CO_3$ .

16 Assuming the atomic weightof a metal M to be 56, find the empirical formula of its oxide containing 70.00% of M.

<b>A16</b>	Element	Percentage composition	Atomic ratio	Least ratio	Whole number ratio
	Metal	70.00	$\frac{70}{56} = 1.25$	$\frac{1.25}{1.25} = 1$	2
	Oxygen	30.00	$\frac{30}{16} = 1.875$	$\frac{1.875}{1.25} = 1.5$	3

Hence, the empirical formula is  $M_2O_3$ .

# STOICHIOMETRY AND STOICHIOMETRIC CALCULTIONS

- The word **stoichiometry** is taken from two Greek words — **stoicheion** (which means element) and **metron** (meaning measurement).
- It deals with the calculation of moles, molecules masses and sometimes volumes of the reactants and products involved in a **balanced chemical equation**.
- The coefficients of the balanced chemical equations are called **stoichiometric coefficients**.
- Stoichiometric coefficients represent the number of moles and molecules of reactants and products in a balanced chemical equations.

# Concept Ladder



Stoichiometry uses all the laws of chemical combination during calculations.

	of 5 mole of ca (1) 22.4 litre (3) 5 × 22.4 litr	ılcium carbonate e	at STP? (2) 2 × 22.4 (4) 3 × 22.4	litre litre	
417	(3) CaCO <sub>3</sub> (s) ———	→ CaO(s) + CO <sub>2</sub> (g	;)		
	1 mole 1 mol CaCO <sub>3</sub> : 5 mol CaCO <sub>3</sub> : At STP 5 mole	1 mole 1 r 1 mol CO <sub>2</sub> 5 mol CO <sub>2</sub> CO <sub>2</sub> will have a v	nole olume of 5 × 22.	4 litre	
	Find out the m	ass of CaO, obta	ined from the t	hermal decomposition	ı of
	(1) 5.6 gm	(2) 2.8 gm	(3) 3.6 gm	(4) 8.6 gm	
418	(2) CaCO <sub>3</sub> (s)	→ CaO(s) + CO <sub>2</sub> (g	;)		
	1 mol CaCO <sub>3</sub> : 1 100 g CaCO <sub>3</sub> : 5	mol CaO 66 g CaO			
	$1 \text{ g CaCO}_3 : \frac{56}{10}$	<mark>6</mark> g CaO			
	$5 g CaCO_3 : \frac{50}{100} = 2.0$	-×5 g CaO ) 8 gm			

Find out the volume of CO<sub>2</sub> produced by the thermal decomposition

5 gm

# Find out the volume of CO<sub>2</sub> obtained at NTP by the thermal decomposition 5gm of CaCO<sub>3</sub>? (1) 22.4 litre (2) 2.245 litre (3) 33.6 litre (4) 1.12 litre A19<sup>(4)</sup> $CaCO_3(s) \longrightarrow CaO(s) + CO_2(g)$ $n_{CaCO_3} = \frac{5}{100}$ $n_{CO_2} = n_{CaCO_3} = \frac{5}{100} = \frac{1}{20}$ $V_{CO_2} = \frac{1}{20} \times V_m = \frac{22.4}{20} = 1.12$ Ltr Find out the volume of $O_2$ obtained from the thermal decomposition of 0.1 mole of Potassium chlorate (KClO<sub>3</sub>) at NTP ? (1) 1.12 Litre (2) 2.24 Litre (3) 3.36 Litre (4) 4.48 Litre A20<sup>(3)</sup> $2\text{KClO}_3 \longrightarrow 2\text{KCl} + 3\text{O}_2$

2 moles of KClO<sub>3</sub> produces 3 moles of O<sub>2</sub> according to the above chemical reaction. So, 1 mol KClO<sub>3</sub> forms 3/2 mol O<sub>2</sub>.  $\therefore$  0.1 mole KClO<sub>3</sub> will give = 0.3/2 × 22.4 litre of O<sub>2</sub> = 3.36 litre

Find out the mass (1) 70 kg (3) 100 kg	of O <sub>2</sub> obtained from 90 kg of water? (2) 80 kg (4) 50 kg
 (0)	

A21 (2)  $2H_2O \longrightarrow 2H_2 + O_2$ 2 mole  $H_2O$  forms 1 mole of  $O_2$ 2 × 18 g of  $H_2O$  forms 32 g of  $O_2$  from above reaction. 1 kg of  $H_2O$  will form  $\frac{32}{2 \times 18}$ kg of  $O_2$ So, from 90 kg of  $H_2O = \frac{32 \times 90}{2 \times 18}$ kg of  $O_2 = 80$  kg of  $O_2$ 



<b>Q23</b> Find out action o (1) 21 gm (3) 63 gn	the weight of iron which w f 18 g of steam? n	vill be converte (2) 42 gm (4) 84 gm	ed into	its oxide (Fe $_{3}O_{4}$ ) by the
A23 (2) From Mola Mass	the compound given the r ratio ratio	following can k 3 mol Fe 3 × 56 g Fe	be ded : :	uced 4 mol H <sub>2</sub> O 4 × 18 g H <sub>2</sub> O

Using stoichiometry  $m_{Fe} = \frac{3 \times 56 \times 18}{4 \times 18} = 42 \text{ gram}$ 

224 Calculate the amount of water (g) produced by the combustion of 16g of methane.

 $A24 \qquad CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(g)$ 

- (i) 16 g of Methane is equal to one mole.
- (ii) From above equation, a mol of  $CH_4(g)$  gives 2 mole of  $H_2O(g)$ .

So, mass of  $H_2O = 2 \times 18 = 36g$ 

# Ideal gas equation :

 PV = nRT P = Pressure of the gasV = Volume of the gas n = mol of gas R = universal gas constant T = temperature in K V<sub>gas</sub> = V<sub>vesel</sub> V<sub>gas</sub> = Free available space for motion  $R = \frac{PV}{nT}$ R = universal gas constant  $R = \frac{0.0821 \text{ atm L}}{\text{mol K}}$ or  $R = \frac{1.99 \text{ Cal}}{\text{mol K}}$  $R = \frac{8.314 \text{ Joule}}{\text{mol K}}$ or PV = nRT at constant P, T  $\left|\frac{V_1}{V_2} = \frac{n_1}{n_2}\right|$  $\uparrow \boxed{V \propto n} \uparrow$ 

• STP / NTP {Standard temp. & Pressure}

# Important points Conversion Factor for gaseous properties : Pressure 1 atm = 760 mm = 760 torr = 1.01325 bar = 101325 Pa 1 bar = 10<sup>5</sup> Pa Volume 1 m<sup>3</sup> = 1000 L 1 dm<sup>3</sup> = 1 L 1 cm<sup>3</sup> = 1 cc = 1 ml Temperature $T_{Ketvin} = 273 + T_{oc}$ $T_{Fahrenheit} = \frac{9}{5}T_{oc} + 32$ 1 L atom = 101.325 J

Pressure}

25 Find out the molar volume of an ideal gas at STP/ NTP?

A25 
$$PV = nRT$$
  $V = \frac{nRT}{P}$ 

At STP: P = 1 Bar, T = 273.15 K

For n = 1 mole and the value of R =  $\frac{0.0821 \text{ atm L}}{\text{mol K}}$ Using ideal gas equation  $V = \frac{\text{nRT}}{\text{P}} = \frac{1 \times 0.0821 \times 273.51}{0.987} = 22.7 \text{ L}$ 

Volume of 1 mol of an ideal gas at 1 atm = 22.4 L

# **Limiting Reagent**

• If the reactants are not taken in the stoichiometric ratios then the reactant which is less than the required amount determines how much product will be formed.

# **Excess Reagent**

- The substance which does not get consumed completely is known as excess reagent.
- The reactant present in excess is called the Excess Reagent.
- **Ex:** If we burn carbon in air (which has an infinite supply of oxygen) then the amount of CO<sub>2</sub> being produced will be governed by the amount of carbon taken. In this case, Carbon is the LR and O<sub>2</sub> is the ER.

# Definition

During a chemical reaction a substance gets consumed completely. This substance is known as limiting reagent.



026 The number o	f litres of air required	to burn 8	3 litres	of $C_2H_2$ is	approximately?
(1) 40		(2) 60			
(3) 80		(4) 100			

A26 For a chemical reaction of the type given below : Given mole at t = 0

 $1 A + 2 B \longrightarrow C + 2 D$ 5 mole 12 mole 0 0

At the completion of reaction the following can be seen, when 12 moles of B reacts with 5 moles of A.

In this case, A gets consumed completely so it behaves as limiting reagent for the reaction

$$C_{2}H_{2} + \frac{5}{2}O_{2} \longrightarrow 2CO_{2} + H_{2}O$$

$$1 \text{ ml}: \frac{5}{2}\text{ ml}$$

$$8 \text{ ml}: \frac{8 \times 5}{2}\text{ ml}$$

$$8 \text{ ml}: 20 \text{ ml of } O_{2}$$

$$20 \times 5 \approx V_{air}$$

$$V_{air} = 100 \text{ ml}$$

# PERCENT YIELD

- The amount of product formed by a chemical reaction is less than the amount predicted by theoretical calculations.
- The ratio of the amount of product formed to the amount predicted when multiplied by 100 gives the percentage yield.

# **Rack your Brain**

Why experimental yield is always less than theoretical or calculated yield?

Percentage Yield =  $\frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100$ 

Q27 For the reaction, CaO + 2HCl → CaCl<sub>2</sub> + H<sub>2</sub>O.
 1.12 gram of CaO si reacted with excess of hydrochloric acid and 1.85 gm CaCl<sub>2</sub> is formed. What is the % yield of the reaction?

A27 CaO + 2HCl  $\longrightarrow$  CaCl<sub>2</sub> + H<sub>2</sub>O 56 gm CaO will produce 111 gm CaCl<sub>2</sub> 1.12 gram of CaO will produce  $\rightarrow \frac{111}{56} \times 1.12 = 2.22$  gm Thus Theoretical yield = 2.22 gm Actual yield = 1.85 gm % yield =  $\frac{1.85}{2.22} \times 100 = 83.33\%$ 

# **PERCENTAGE PURITY**

• Depending upon the mass of product, the equivalent amount of reactant present is determined with the help of given chemical equation. When we known the actual amount of reactant taken and the amount calculated with the help of a chemical equation, the purity can be determined.





28 Calcualte the amount of (CaO) in kg that can be produced by heating 200 kg lime stone taht is 90% pure caCO<sub>3</sub>.

A28 Mass of Pure  $CaCO_3 = \frac{200 \times 90}{100} = 180 \text{ kg}$   $CaCO_3 \longrightarrow CaO + CO_2$   $100 \text{ kg} \qquad 56 \text{ kg}$   $180 \qquad x$  $\frac{100}{180} = \frac{56}{x} \Rightarrow x = 100.8 \text{ kg}$ 

# Eudiometry

- It is the special part of stoichiometry.
- In this, we deal with absorption of gases during a chemical process.

# Some Absorbents of Gases

The absorbent which is used for specific gas is listed below:

# Definition

The process of determining the constituents of a gaseous mixture by means of the eudiometer, or for ascertainnig the purity of the air or the amount of oxygen in it.

Absorbent	Gas or gases absorbed
Turpentine oil	0 <sub>3</sub>
Alkaline pyrogallol	0 <sub>2</sub>
Ferrous sulphate solution	NO
Heated magnesium	N <sub>2</sub>
Heated palladium	H <sub>2</sub>
Ammonical coprous chloride	O <sub>2</sub> , CO, C <sub>2</sub> H <sub>2</sub>
Copper sulphate solution	H <sub>2</sub> S, PH <sub>3</sub> , AsH <sub>3</sub>
Conc. H <sub>2</sub> SO <sub>4</sub>	H <sub>2</sub> O i.e., moisture, NH <sub>3</sub>
NaOH or KOH solution	CO <sub>2</sub> , NO <sub>2</sub> , SO <sub>2</sub> , X <sub>2</sub> , all acidic oxides

# 90 ml of pure dry O<sub>2</sub> is subjected to electric discharge, if only 10 % of it is converted into O<sub>3</sub>, volume of the mixture of gases (O<sub>2</sub> & O<sub>3</sub>) after the reaction will be \_\_\_\_\_\_and after passing through turpentine oil will be \_\_\_\_\_\_.

(1) 8	4 ml,	ml, 78 ml
(3) 7	8 ml,	ml, 84 ml

(2) 81 ml, 87 ml (4) 87 ml, 81 ml

# A29

 $\begin{array}{cccc} & & & & & & & & & & \\ \text{Initially} & & & & & & & & & \\ \text{After reaction} & & & & & & & & & \\ \text{After reaction} & & & & & & & & & & \\ \text{Mix} = 81 \text{ ml } \text{O}_2 + 6 \text{ ml } \text{O}_3 = 87 \text{ ml} \\ \text{Turpentine oil (It absorbes } \text{O}_3 \text{ gas)} \\ \text{V}_{\text{remaining}} = 81 \text{ ml of } \text{O}_2 \end{array}$ 

# **CONCENTRATION TERMS**

- A solution is defined as the homogeneous mixture of two or more substances, composition of which may vary within the limits.
- Solution is a special type of mixture in which substances are intermixed so intimately that they cannot be observed as separated components.
- The substance which is dissolved is called solute while medium in which solute is dissovled to get a homogeneous mixture is called the solvent.
- A solution is termed as binary and ternary if it consists of two and three components





How does strength of solution change by increase in temperature?



respectively.

# Percentage

- It refers tot he amount of the solute per 100 parts of the solution. It can also be called as parts per hundred (pph). It can be expressed be any of following four methods:
- (i) Weight by weight percentage (%w/w)

 $= \frac{\text{Wt. of solute(g)}}{\text{Wt. of solution(g)}} \times 100$ 

e.g., 10% Na<sub>2</sub>CO<sub>3</sub> solution w/w means 10g of Na<sub>2</sub>CO<sub>3</sub> is dissolved in 100 g of the solution.

# (ii) Weight by volume percent (%w/v)

 $= \frac{\text{Wt. of solute(g)}}{\text{Volume of solution(cm}^3)} \times 100$ 

e.g., 10%  $Na_2CO_3$  (w/v) means 10 g  $Na_2CO_3$  is dissolved in 100 cm<sub>3</sub> of solution.

# (iii) Volume by bolume percent (%v/v)

 $= \frac{\text{Volume of solute(cm}^3)}{\text{Volume of solution(cm}^3)} \times 100$ 

e.g., 10% ethanol (v/v) means 10 cm<sup>3</sup> of ethanol dissolved in 100 cm<sup>3</sup> of solution.

# (iv) Volume by weight percent (%v/w)

 $= \frac{\text{Volume of solute(cm^3)}}{\text{Wt. of solution(g)}} \times 100$ 

e.g, 10% ethanol (v/w) means 10cm<sup>3</sup> of ethanol dissolved in 100g of solution.







Is there any effect of change in temperature on %v/v?



35.

# Parts per million (ppm) and parts per billion (ppb)

- When a solute is present in very small quantity, it is convenient to express the concentration in parts per million and parts per billion.
- It is the number of parts of solute per million (10<sup>6</sup>) or per billion (10<sup>9</sup>) parts of solution.
- It is independent of the temperature.

 $ppm = \frac{Mass of solute}{Total mass of solution} \times 10^{6}$  $ppb = \frac{Mass of solute}{Total mass of solution} \times 10^{9}$ 

#### **Rack your Brain**



What is the application of ppm, ppb or ppt in daily life?

230 Calculate the parts per million of SO<sub>2</sub> gas in 250 mL water containing 5 × 10<sup>04</sup> g of SO<sub>2</sub> gas.

A30 Mass of SO<sub>2</sub> gas = 5 × 10<sup>-4</sup>g; Mass of H<sub>2</sub>O = Volume × Density = 250 cm<sup>3</sup> × 1 g cm<sup>-3</sup> = 250 g  $\therefore$  Parts per million of SO<sub>2</sub> gas =  $\frac{5 \times 10^{-4}}{250 \text{ g}} \times 10^{-6} = 2$ 

# Molarity (M)

- Molarity is defined as the number of moles of the solute per liter of solution. Unit of molarity is mol/dm<sup>3</sup> or mol/liter.
- For example, one molar (1 M) solution of sugar means the solution contains 1 mole of sugar per litre of the solution. Solution in terms of molarity is generally expreesd as,
- Mathematically, molarity can be calculated by following formulas:



(i)  $M = \frac{\text{No. of moles of solute(n)}}{\text{Vol. of solution in litres}} = \frac{\text{Wt. of solute(gm)}}{\text{gm mol.wt. of solute}} \times \frac{1000}{\text{wt. of solution(ml)}}$ 

(ii) If molarity and volume of the solution are changed from M<sub>1</sub>, V<sub>1</sub> to M<sub>2</sub>, V<sub>2</sub>.

Then,  $M_1V_1 = M_2V_2$ 

(iii) (iii)In the balanced chemical equation, if n1 moles of reactant-1 react with n2 moles of reactant-2. Then,

$$n_1 A + n_2 B \rightarrow Product$$
  
 $M_1 V_1 = M_2 V_2$ 

$$\frac{1}{n_1} = \frac{1}{n_2}$$

(iv) (iv) If two solutions of same solute are mixed then molarity (M) of resulting solution

$$\mathsf{M} = \frac{\mathsf{M}_1 \mathsf{V}_1 + \mathsf{M}_2 \mathsf{V}_2}{\left(\mathsf{V}_1 + \mathsf{V}_2\right)}$$

A bottle of commercial sulphuric acid (density 1.787 g ml-1) is labelled as 86% by weight. What is the molarity of acid?

Before

High Concentration Of Solute

A31 Molarity of  $H_2SO_4 = \frac{Wt. \text{ of } H_2SO_4 \text{ in 1L solution}}{\text{mol. wt. of } H_2SO_4}$ But wt. of given  $H_2SO_4$  per litre  $= \frac{86}{100} \times 1.787 \times 1000 = 1536.82 \text{ g}$ Hence molarity of  $H_2SO_4 = \frac{1536.82}{98} = 15.68 \text{ mol } L^{-1}$ 

Q32 A sample contains  $I_2$  and benzene. The mole fraction of  $I_2$  = 0.2. Calculate molarity of solution if

- (i) density of solution is d gm/ml
- (ii) density of  $\rm I_{_2}$  & benzene are  $\rm d_{_{12}}$  &  $\rm d_{_{benzene}}$

(ii) M = 
$$\frac{0.2}{\left(\frac{50.8}{d_{l_2}} + \frac{62.4}{d_{benzene}}\right)} \times 1000$$







# Formality (F)

• Formality of solution may be defined as the number of gram formula unis of the ionic solute dissolved per litre of the solution.

Formality (F) = 
$$\frac{\text{Number of gram formula units of solute}}{\text{Volume of solution in litres}}$$
$$= \frac{\text{Mass of ionic solute (g)}}{\text{gram formula unit mass of solute × Volume of solution (l)}}$$

What will be the formality of KNO<sub>3</sub> solution having strength equal to 2.02 g per litre?

 $\Delta$  33 Sterngth of KNO<sub>3</sub> = 2.02 gL<sup>-1</sup>

:. Formality of KNO<sub>3</sub> = 
$$\frac{\text{strength in gL}^{-1}}{\text{g-formula wt. of KNO}_3} = \frac{2.02}{101} = 0.02 \text{ F}$$

# Molality (m)

- It is the number of moles of the solute per 1000 g of the solvent.
- Unit of molarity is mol/kg.
- Mathematically, molality can be calculated by the following formulae,

(i) 
$$m = \frac{\text{Number of moles of solute}}{\text{Weight of solvent in kg}} = \frac{\text{Number of moles of solute}}{\text{Weight of solvent in gm}} \times 1000$$

(ii) 
$$m = \frac{Wt. of solute}{Mol. wt. of solute} \times \frac{1000}{Weight of solvent in gm}$$

Molality (m) = 
$$\frac{M \times 1000}{1000d - MM_{o}}$$

24 Calculate the Molarity and molality of a 98% by mass of H<sub>2</sub>SO<sub>4</sub> solution having a density of 1.25 g/cc.

A34 
$$H_2SO_4$$
 taken = 98%  
100g of solution contains 98g  $H_2SO_4$ .  
mass of solute,  $H_2SO_4$  = 98g  
mass of volvent = 100 - 98 = 2g = 0.002 kg  
moles of solute,  $H_2SO_4 = \frac{98}{98} = 1$   
Volume of solution =  $\frac{\text{mass of solution}}{\text{density}} = \frac{100}{1.25} = 80 \text{ mL} = 0.08 \text{ L}$   
Molarity, M =  $\frac{\text{moles of solute}}{\text{volume of solution}(L)} = \frac{1}{0.08} = 12.5 \text{ M}$   
Molarity, M =  $\frac{\text{moles of solute}}{\text{mass of solute}} = \frac{1}{0.02} = 500 \text{ m}$ 

# Mole fraction (X)

- It is defined as ratio of number of moles of a component to total moles of all components (solvent and solute) present in solution.
- It is denoted by the letter X. Number of moles of component A is given by,  $n_A = \frac{W_A}{M_A}$

Number of moles of component B is given by,  $n_{_B} = \frac{W_{_B}}{M_{_B}}$ 

Total number of moles of A and B =  $n_A + n_B$ Mole fraction of A,  $X_A = \frac{n_A}{n_A + n_B}$ 

Mole fraction of B,  $X_{_B} = \frac{n_{_B}}{n_{_A} + n_{_B}}$ 

The sum of mole fractions of all the components in the solution is always one.

$$X_{_{A}} + X_{_{B}} = \frac{n_{_{A}}}{n_{_{A}} + n_{_{B}}} + \frac{n_{_{B}}}{n_{_{A}} + n_{_{B}}} = 1$$

Concept LadderImage: Relationship between molality, moles and molefraction.Image: molefraction.Image: molefraction.Image: molefraction moles of moles of solute not molefraction of solute xolute not molefraction of solute not molefraction not molefracti

Mole Concept

Find out the masses of acid and water requried to prepare 1 mle of CH<sub>3</sub>COOH solution of 0.3 mole fraction of CH<sub>3</sub>COOH.

A35  $X_{CH_3COOH} = 0.3$   $X_{H_2O} = 1 - 0.3 = 0.7$ Wt. of CH<sub>3</sub>COOH =  $X_{CH_3COOH} \times mol. wt.(CH_3COOH) = 0.3 \times 60 = 18 g$ Wt. of water =  $X_{H_2O} \times mol. wt.(H_2O) = 0.7 \times 18 = 12.6 g$ 

**36** From 160 g of SO<sub>2</sub> (g) sample, 1.2046 x 10<sup>24</sup> molecules of SO<sub>2</sub> are removed then find out the volume of left over SO<sub>2</sub> (g) at STP.

A36 Given moles =  $\frac{160}{64}$  = 2.5 Removed moles =  $\frac{1.2046 \times 10^{24}}{6.023 \times 10^{23}}$  = 2

> Number of moles left = 0.5. Remaining volume at STP =  $0.5 \times 22.4 = 11.2$  lit..

When x gram of a certain metal brunt in 1.5 g oxygen to give 3.0 g of its oxide.
1.20 g of the same metal heated in a steam gave 2.40 g of its oxide. shows the these result illustrate the law of constant or definite proportion

A37 weight of metal = 3.0 – 1.5 = 1.5 g So, weight of metal : weight of oxygen = 1.5 : 1.5 = 1 : 1 Similarly, in second case, weight of oxygen = 2.4 – 1.2 = 1.2 g so weight of metal : weight of oxygen = 1.2 : 1.2 = 1 : 1 so, them results illustrate law of constant proportion.



# **Chapter Summary**

- Atom is the fundamental unit of matter which is further indivisible i.e. atom can ٠ neither be created nor be destroyed.
- Actual mass of the mass of one atom or one molecule of a substance is called as actual mass.
- 1 amu =  $\frac{1}{12}$  × mass of one C-12 atom = 1.66 × 10<sup>-24</sup> g or 1.66 × 10<sup>-27</sup> kg 1 mole of atoms is also termed as 1 gm-atom, 1 mole of ions is termed as 1 gm-٠ ion and 1 mole of molecule termed as 1 gm - molecule
- Relation between the molecular formula and Empirical formula
- Molecular mass n =

Empirical Formula mass

Vapour density = molecular mass ٠

- **Limiting reagent :** Calculating amount of anyone product obtained taking each ٠ reactant one by one irrespective of other reactants. The one giving least product is limiting reagent.
- For reversible reaction, the actual amount of any limiting reagent consumed in ٠ such incomplete reaction is given by [% yield × given mole of limit reagent]
- Measuring volume is equivalent to counting number of molecules of gas General Concentration term :

(a) Density = <u>mass</u>, Unit : gm/cc volume Density of any substance (b) Relative density =

Density of reference substance

(c) Specific gravity = <u>Density of any substance</u> Density of water at 4°C

Density of vapour at some temperature and pressure (d) Vapour density = Density of H<sub>g</sub>as at same temperature and pressure

**Concentration Terms** 

$$\% = \left(\frac{v}{V}\right) = \frac{\text{volume of solute}}{\text{volume solution}} \times 100$$
  
Mole % =  $\frac{\text{Moles of solute}}{\text{Total moles}} \times 100$ 

